



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





DCAMKL3 siRNA (m): sc-142890

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including cell division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine (Ser/Thr) protein kinases. DCAMKL3 (doublecortin-like and CAM kinase-like 3), also known as DCLK3 or DCDC3C, is a 648 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one protein kinase domain. One of several members of the Ser/Thr protein kinase family, DCAMKL3 functions to catalyze the ATP-dependent attachment of a phospho residue to target proteins, an event that may play a role in various cellular processes.

REFERENCES

1. Bairoch, A., et al. 1988. Sequence patterns in protein kinases. *Nature* 331: 22.
2. Hanks, S.K., et al. 1988. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241: 42-52.
3. Hanks, S.K., et al. 1991. Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Meth. Enzymol.* 200: 38-62.
4. Sapir, T., et al. 2000. Doublecortin mutations cluster in evolutionarily conserved functional domains. *Hum. Mol. Genet.* 9: 703-712.
5. Reiner, O., et al. 2006. The evolving doublecortin (DCX) superfamily. *BMC Genomics* 7: 188.
6. Coquelle, F.M., et al. 2006. Common and divergent roles for members of the mouse DCX superfamily. *Cell Cycle* 5: 976-983.
7. Cierpicki, T., et al. 2006. The DC-module of doublecortin: dynamics, domain boundaries, and functional implications. *Proteins* 64: 874-882.

CHROMOSOMAL LOCATION

Genetic locus: Dclk3 (mouse) mapping to 9 F3.

PRODUCT

DCAMKL3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DCAMKL3 shRNA Plasmid (m): sc-142890-SH and DCAMKL3 shRNA (m) Lentiviral Particles: sc-142890-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DCAMKL3 siRNA (m) is recommended for the inhibition of DCAMKL3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DCAMKL3 gene expression knockdown using RT-PCR Primer: DCAMKL3 (m)-PR: sc-142890-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.