



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AWAT1 siRNA (m): sc-143019

BACKGROUND

AWAT1 (Acyl-CoA wax alcohol acyltransferase 1) is a 328 amino acid protein that belongs to the diacylglycerol acyltransferase family. AWAT1 localizes to endoplasmic reticulum where it is considered a multi-pass membrane protein. AWAT1 is an acyltransferase that predominantly esterifies long chain (wax) alcohols with acyl-CoA-derived fatty acids to produce wax esters, suggesting that it plays a central role in lipid metabolism in skin. Predominantly expressed in skin, where it is limited to sebaceous gland, AWAT1 is expressed in more mature, centrally located cells prior to rupture and sebum release. AWAT1 is also ubiquitously expressed, except in spleen, with higher levels in thymus, prostate and testis. The AWAT1 gene is conserved in chimpanzee, canine, bovine, mouse and rat, and maps to human chromosome X.

REFERENCES

1. Strausberg, R.L., Feingold, E.A., Grouse, L.H., Derge, J.G., Klausner, R.D., Collins, F.S., Wagner, L., Shenmen, C.M., Schuler, G.D., Altschul, S.F., Zeeberg, B., Buetow, K.H., Schaefer, C.F., Bhat, N.K., Hopkins, R.F., et al. 2002. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. *Proc. Natl. Acad. Sci. USA* 99: 16899-16903.
2. Winter, A., van Eckeveld, M., Bininda-Emonds, O.R., Habermann, F.A. and Fries, R. 2003. Genomic organization of the DGAT2/MOGAT gene family in cattle (*Bos taurus*) and other mammals. *Cytogenet. Genome Res.* 102: 42-47.
3. Orland, M.D., Anwar, K., Cromley, D., Chu, C.H., Chen, L., Billheimer, J.T., Hussain, M.M. and Cheng, D. 2005. Acyl coenzyme A dependent retinol esterification by acyl coenzyme A: diacylglycerol acyltransferase 1. *Biochim. Biophys. Acta* 1737: 76-82.
4. Turkish, A.R., Henneberry, A.L., Cromley, D., Padamsee, M., Oelkers, P., Bazzi, H., Christiano, A.M., Billheimer, J.T. and Sturley, S.L. 2005. Identification of two novel human acyl-CoA wax alcohol acyltransferases: members of the diacylglycerol acyltransferase 2 (DGAT2) gene superfamily. *J. Biol. Chem.* 280: 14755-14764.
5. Ross, M.T., Grafham, D.V., Coffey, A.J., Scherer, S., McLay, K., Muzny, D., Platzer, M., Howell, G.R., Burrows, C., Bird, C.P., Frankish, A., Lovell, F.L., Howe, K.L., Ashurst, J.L., Fulton, R.S., Sudbrak, R., Wen, G., et al. 2005. The DNA sequence of the human X chromosome. *Nature* 434: 325-337.
6. Turkish, A. and Sturley, S.L. 2007. Regulation of triglyceride metabolism. I. Eukaryotic neutral lipid synthesis: "many ways to skin ACAT or a DGAT". *Am. J. Physiol. Gastrointest. Liver Physiol.* 292: G953-G957.
7. Holmes, R.S. 2010. Comparative genomics and proteomics of vertebrate diacylglycerol acyltransferase (DGAT), acyl CoA wax alcohol acyltransferase (AWAT) and monoacylglycerol acyltransferase (MGAT). *Comp. Biochem. Physiol. Part D Genomics Proteomics* 5: 45-54.
8. SWISS-PROT/TrEMBL (Q58HT5). World Wide Web URL: <http://www.uniprot.org/uniprot/Q58HT5>

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Awat1 (mouse) mapping to X C3.

PRODUCT

AWAT1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AWAT1 shRNA Plasmid (m): sc-143019-SH and AWAT1 shRNA (m) Lentiviral Particles: sc-143019-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AWAT1 siRNA (m) is recommended for the inhibition of AWAT1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AWAT1 gene expression knockdown using RT-PCR Primer: AWAT1 (m)-PR: sc-143019-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.