



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DNAH3 siRNA (m): sc-143079

BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Cytoplasmic or axonemal Dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; the complex transports cellular cargos towards the central region of the cell. Axonemal dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell. DNAH3 (dynein, axonemal, heavy chain 3), also known as DNAHC3B, DLP3 or FLJ31947, is a 4,116 amino acid member of the dynein heavy chain protein family. Expressed primarily in testis and trachea, DNAH3 contains twenty-one LRR repeats. DNAH3 is a force generating protein of respiratory cilia, and is thought to be involved in sperm motility through sperm flagellar assembly.

REFERENCES

1. Chapelin, C., Duriez, B., Magnino, F., Goossens, M., Escudier, E. and Amselem, S. 1997. Isolation of several human axonemal dynein heavy chain genes: genomic structure of the catalytic site, phylogenetic analysis and chromosomal assignment. *FEBS Lett.* 412: 325-330.
2. Neesen, J., Koehler, M.R., Kirschner, R., Steinlein, C., Kreutzberger, J., Engel, W. and Schmid, M. 1997. Identification of dynein heavy chain genes expressed in human and mouse testis: chromosomal localization of an axonemal dynein gene. *Gene* 200: 193-202.
3. Milisav, I. 1998. Dynein and dynein-related genes. *Cell Motil. Cytoskeleton* 39: 261-272.
4. Milisav, I. and Affara, N.A. 1998. A potential human axonemal dynein heavy-chain gene maps to 17q25. *Mamm. Genome* 9: 404-407.
5. Maiti, A.K., Mattei, M.G., Jorissen, M., Volz, A., Zeigler, A. and Bouvagnet, P. 2000. Identification, tissue specific expression, and chromosomal localisation of several human dynein heavy chain genes. *Eur. J. Hum. Genet.* 8: 923-932.
6. Carson, J.L., Reed, W., Lucier, T., Brighton, L., Gambling, T.M., Huang, C.H. and Collier, A.M. 2002. Axonemal dynein expression in human fetal tracheal epithelium. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 282: L421-L430.
7. Seetharam, R.N. and Satir, P. 2005. High speed sliding of axonemal microtubules produced by outer arm dynein. *Cell Motil. Cytoskeleton* 60: 96-103.

CHROMOSOMAL LOCATION

Genetic locus: Dnah3 (mouse) mapping to 7 F2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

DNAH3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DNAH3 shRNA Plasmid (m): sc-143079-SH and DNAH3 shRNA (m) Lentiviral Particles: sc-143079-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DNAH3 siRNA (m) is recommended for the inhibition of DNAH3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DNAH3 gene expression knockdown using RT-PCR Primer: DNAH3 (m)-PR: sc-143079-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.