



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DnaJC25 siRNA (m): sc-143106

BACKGROUND

With the presence of the J domain defining a protein as a member, the DnaJ family has evolved with diverse cellular localization and functions and is one of the largest chaperone families. DnaJ heat-shock-induced proteins are derived from the bacterium *Escherichia coli* and are controlled by the htpR regulatory protein. DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis. Members of this family contain cysteine-rich regions composed of zinc fingers that form a peptide-binding domain responsible for chaperone function. DnaJ proteins are important mediators of proteolysis and are involved in the regulation of protein degradation, exocytosis and endocytosis. DnaJC25 (DnaJ homolog subfamily C member 25) is a 360 amino acid multi-pass membrane protein that contains one J domain and is expressed as three isoforms produced by alternative splicing.

REFERENCES

1. Saito, H. and Uchida, H. 1978. Organization and expression of the dnaJ and dnaK genes of *Escherichia coli* K12. Mol. Gen. Genet. 164: 1-8.
2. Georgopoulos, C.P., Lundquist-Heil, A., Yochem, J. and Feiss, M. 1980. Identification of the *E. coli* dnaJ gene product. Mol. Gen. Genet. 178: 583-588.
3. Suh, W.C., Burkholder, W.F., Lu, C.Z., Zhao, X., Gottesman, M.E. and Gross, C.A. 1998. Interaction of the Hsp70 molecular chaperone, DnaK, with its cochaperone DnaJ. Proc. Natl. Acad. Sci. USA 95: 15223-15228.
4. Tomoyasu, T., Ogura, T., Tatsuta, T. and Bukau, B. 1998. Levels of DnaK and DnaJ provide tight control of heat shock gene expression and protein repair in *Escherichia coli*. Mol. Microbiol. 30: 567-581.
5. Stewart, G.R., Robertson, B.D. and Young, D.B. 2004. Analysis of the function of mycobacterial DnaJ proteins by overexpression and microarray profiling. Tuberculosis 84: 180-187.
6. Shi, Y.Y., Hong, X.G. and Wang, C.C. 2005. The C-terminal (331-376) sequence of *Escherichia coli* DnaJ is essential for dimerization and chaperone activity: a small angle X-ray scattering study in solution. J. Biol. Chem. 280: 22761-22768.
7. Robichon, C., Varret, M., Le Liepvre, X., Lasnier, F., Hajdich, E., Ferre, P. and Dugail, I. 2006. DnaJA4 is a SREBP-regulated chaperone involved in the cholesterol biosynthesis pathway. Biochim. Biophys. Acta 1761: 1107-1113.
8. Acebrón, S.P., Fernández-Sáiz, V., Taneva, S.G., Moro, F. and Muga, A. 2008. DnaJ recruits DnaK to protein aggregates. J. Biol. Chem. 283: 1381-1390.

CHROMOSOMAL LOCATION

Genetic locus: Dnajc25 (mouse) mapping to 4 B3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

DnaJC25 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DnaJC25 shRNA Plasmid (m): sc-143106-SH and DnaJC25 shRNA (m) Lentiviral Particles: sc-143106-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DnaJC25 siRNA (m) is recommended for the inhibition of DnaJC25 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DnaJC25 gene expression knockdown using RT-PCR Primer: DnaJC25 (m)-PR: sc-143106-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.