



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



DSN1 siRNA (m): sc-143178

BACKGROUND

DSN1, also known as MIS13, is a 356 amino acid protein that is associated with the kinetochore and functions as a component of the MIS12 protein complex that works to ensure proper kinetochore formation and spindle checkpoint activity. Expressed as multiple alternatively spliced isoforms, DSN1 is also required for correct chromosome alignment during metaphase. The gene encoding DSN1 maps to human chromosome 20. Comprising approximately 2% of the human genome, chromosome 20 contains nearly 63 million bases that encode over 600 genes, some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 609175. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Pinsky, B.A., et al. 2003. An MTW1 complex promotes kinetochore biorientation that is monitored by the iPL1/Aurora protein kinase. *Dev. Cell* 5: 735-745.
3. Cheeseman, I.M., et al. 2004. A conserved protein network controls assembly of the outer kinetochore and its ability to sustain tension. *Genes Dev.* 18: 2255-2268.
4. Obuse, C., et al. 2004. A conserved MIS12 centromere complex is linked to heterochromatic HP1 and outer kinetochore protein Zwint-1. *Nat. Cell Biol.* 6: 1135-1141.
5. Kline, S.L., et al. 2006. The human MIS12 complex is required for kinetochore assembly and proper chromosome segregation. *J. Cell Biol.* 173: 9-17.
6. Nousiainen, M., et al. 2006. Phosphoproteome analysis of the human mitotic spindle. *Proc. Natl. Acad. Sci. USA* 103: 5391-5396.
7. Yang, Y., et al. 2008. Phosphorylation of HsMIS13 by Aurora B kinase is essential for assembly of functional kinetochore. *J. Biol. Chem.* 283: 26726-26736.

CHROMOSOMAL LOCATION

Genetic locus: Dsn1 (mouse) mapping to 2 H1.

PRODUCT

DSN1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DSN1 shRNA Plasmid (m): sc-143178-SH and DSN1 shRNA (m) Lentiviral Particles: sc-143178-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DSN1 siRNA (m) is recommended for the inhibition of DSN1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DSN1 gene expression knockdown using RT-PCR Primer: DSN1 (m)-PR: sc-143178-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.