



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



DTX3L siRNA (m): sc-143183

BACKGROUND

The Deltex family is responsible for influencing Notch signaling and may regulate transcription through interactions with specific transcription factors. Deltex proteins have a basic N-terminus, a central proline-rich region and a C-terminal RING-type zinc finger domain, a motif often found in ubiquitin-protein isopeptide ligases (E3). The RING-type zinc finger domain binds two Zn²⁺ atoms and forms a cross-brace motif that is essential for many proteins involved in the ubiquitination pathway. DTX3L (Deltex-3-like), also known as BBAP, is a 740 amino acid protein that is similar to Deltex-3 and acts as a ubiquitin ligase *in vitro*. DTX3L can heterodimerize with Deltex-1, a transcriptional regulator, thereby enhancing the activity of the E3 ubiquitin ligase complex and increasing the influence of E3 on the Notch signaling pathway.

REFERENCES

1. Matsuno, K., Eastman, D., Mitsiades, T., Quinn, A.M., Carcanciu, M.L., Ordentlich, P., Kadesch, T. and Artavanis-Tsakonas, S. 1998. Human deltex is a conserved regulator of Notch signalling. *Nat. Genet.* 19: 74-78.
2. Yamamoto, N., Yamamoto, S., Inagaki, F., Kawaichi, M., Fukamizu, A., Kishi, N., Matsuno, K., Nakamura, K., Weinmaster, G., Okano, H. and Nakafuku, M. 2001. Role of Deltex-1 as a transcriptional regulator downstream of the Notch receptor. *J. Biol. Chem.* 276: 45031-45040.
3. Izon, D.J., Aster, J.C., He, Y., Weng, A., Karnell, F.G., Patriub, V., Xu, L., Bakkour, S., Rodriguez, C., Allman, D. and Pear, W.S. 2002. Deltex-1 redirects lymphoid progenitors to the B cell lineage by antagonizing Notch 1. *Immunity* 16: 231-243.
4. Takeyama, K., Aguiar, R.C., Gu, L., He, C., Freeman, G.J., Kutok, J.L., Aster, J.C. and Shipp, M.A. 2003. The BAL-binding protein BBAP and related deltex family members exhibit ubiquitin-protein isopeptide ligase activity. *J. Biol. Chem.* 278: 21930-21937.
5. Cui, X.Y., Hu, Q.D., Tekaya, M., Shimoda, Y., Ang, B.T., Nie, D.Y., Sun, L., Hu, W.P., Karsak, M., Duka, T., Takeda, Y., Ou, L.Y., Dawe, G.S., Yu, F.G., Ahmed, S., Jin, L.H., Schachner, M., Watanabe, K., Arsenijevic, Y. and Xiao, Z.C. 2004. NB-3/Notch 1 pathway via Deltex-1 promotes neural progenitor cell differentiation into oligodendrocytes. *J. Biol. Chem.* 279: 25858-25865.
6. Juszczynski, P., Kutok, J.L., Li, C., Mitra, J., Aguiar, R.C. and Shipp, M.A. 2006. BAL1 and BBAP are regulated by a γ interferon-responsive bidirectional promoter and are overexpressed in diffuse large B-cell lymphomas with a prominent inflammatory infiltrate. *Mol. Cell. Biol.* 26: 5348-5359.

CHROMOSOMAL LOCATION

Genetic locus: Dtx3l (mouse) mapping to 16 B3.

PRODUCT

DTX3L siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DTX3L shRNA Plasmid (m): sc-143183-SH and DTX3L shRNA (m) Lentiviral Particles: sc-143183-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DTX3L siRNA (m) is recommended for the inhibition of DTX3L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DTX3L gene expression knockdown using RT-PCR Primer: DTX3L (m)-PR: sc-143183-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.