



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Dynactin 3 siRNA (m): sc-143203

BACKGROUND

Dynactin, a multisubunit complex, is a cytoplasmic Dynein-interacting protein that functions as the "receptor" for the Dynein microtubule motor. Dynactin/Dynein binding may be required for most, if not all, cytoplasmic Dynein-driven activities and is thought to contribute to the functional diversity of Dynein. Enriched in neurons, Dynactin also binds to microtubules and has been shown to function in diverse processes, including organelle transport, formation of the mitotic spindle and cytokinesis. Dynactin 3, also known as Dynactin complex subunit 22 kDa subunit or p22, is a 186 amino acid subunit of the Dynactin complex. Dynactin 3 localizes to punctate cytoplasmic structures, to the centrosome during interphase and to the kinetochores and spindle poles throughout mitosis. Ubiquitously expressed, Dynactin 3 is found at highest levels in pancreas and muscle.

REFERENCES

1. Karki, S., LaMonte, B. and Holzbaur, E.L. 1998. Characterization of the p22 subunit of dynactin reveals the localization of cytoplasmic dynein and dynactin to the midbody of dividing cells. *J. Cell Biol.* 142: 1023-1034.
2. Karki, S., Tokito, M.K. and Holzbaur, E.L. 2000. A dynactin subunit with a highly conserved cysteine-rich motif interacts directly with Arp1. *J. Biol. Chem.* 275: 4834-4839.
3. Mayor, T., Stierhof, Y.D., Tanaka, K., Fry, A.M. and Nigg, E.A. 2000. The centrosomal protein C-Nap1 is required for cell cycle-regulated centrosome cohesion. *J. Cell Biol.* 151: 837-846.
4. Mills, D.R. and Jackson, C.L. 2001. Assignment of p22 dynactin light chain (DCTN3) to human chromosome region 9p13 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 92: 166.
5. Takahashi, M., Yamagiwa, A., Nishimura, T., Mukai, H. and Ono, Y. 2002. Centrosomal proteins CG-NAP and kendrin provide microtubule nucleation sites by anchoring γ -Tubulin ring complex. *Mol. Biol. Cell* 13: 3235-3245.
6. Casenghi, M., Meraldi, P., Weinhart, U., Duncan, P.I., Körner, R. and Nigg, E.A. 2003. Polo-like kinase 1 regulates Nlp, a centrosome protein involved in microtubule nucleation. *Dev. Cell* 5: 113-125.
7. Lehner, B., Semple, J.I., Brown, S.E., Counsell, D., Campbell, R.D. and Sanderson, C.M. 2004. Analysis of a high-throughput yeast two-hybrid system and its use to predict the function of intracellular proteins encoded within the human MHC class III region. *Genomics* 83: 153-167.
8. Lim, J., Hao, T., Shaw, C., Patel, A.J., Szabó, G., Rual, J.F., Fisk, C.J., Li, N., Smolyar, A., Hill, D.E., Barabási, A.L., Vidal, M. and Zoghbi, H.Y. 2006. A protein-protein interaction network for human inherited ataxias and disorders of Purkinje cell degeneration. *Cell* 125: 801-814.
9. Petretti, C., Savoian, M., Montembault, E., Glover, D.M., Prigent, C. and Giet, R. 2006. The PITSLRE/CDK11p58 protein kinase promotes centrosome maturation and bipolar spindle formation. *EMBO Rep.* 7: 418-424.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Dctn3 (mouse) mapping to 4 A5.

PRODUCT

Dynactin 3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Dynactin 3 shRNA Plasmid (m): sc-143203-SH and Dynactin 3 shRNA (m) Lentiviral Particles: sc-143203-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Dynactin 3 siRNA (m) is recommended for the inhibition of Dynactin 3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Dynactin 3 gene expression knockdown using RT-PCR Primer: Dynactin 3 (m)-PR: sc-143203-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.