



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PAMR1 siRNA (m): sc-143257

BACKGROUND

PAMR1 (peptidase domain containing associated with muscle regeneration 1), also known as RAMP or FP938, is a 720 amino acid secreted protein that belongs to the peptidase S1 family. Consisting of one CUB domain, an EGF-like domain, a peptidase S1 domain and two sushi (CCP/SCR) domains, PAMR1 may participate in regeneration of skeletal muscle. PAMR1 is strongly down-regulated in muscle cell lines derived from Duchenne muscular dystrophy (DMD) patients compared to a normal muscle cell line. DMD is the second most common genetically inherited disease in humans and is characterized by progressive limb-girdle distribution of muscle weakness. PAMR1 exists as two alternatively spliced isoforms and is encoded by a gene located on human chromosome 11, which contains 135 million base pairs and 1,400 genes, making up around 4% of human genomic DNA.

REFERENCES

1. Anderson, J.L., Head, S.I., Rae, C. and Morley, J.W. 2002. Brain function in Duchenne muscular dystrophy. *Brain* 125: 4-13.
2. Nakayama, Y., Nara, N., Kawakita, Y., Takeshima, Y., Arakawa, M., Katoh, M., Morita, S., Iwatsuki, K., Tanaka, K., Okamoto, S., Kitamura, T., Seki, N., Matsuda, R., Matsuo, M., Saito, K. and Hara, T. 2004. Cloning of cDNA encoding a regeneration-associated muscle protease whose expression is attenuated in cell lines derived from Duchenne muscular dystrophy patients. *Am. J. Pathol.* 164: 1773-1782.
3. Hara, T., Nakayama, Y. and Nara, N. 2005. Regenerative medicine of skeletal muscle. *Rinsho Shinkeigaku* 45: 880-882.
4. Taylor, T.D., Noguchi, H., Totoki, Y., Toyoda, A., Kuroki, Y., Dewar, K., Lloyd, C., Itoh, T., Takeda, T., Kim, D.W., She, X., Barlow, K.F., Bloom, T., Bruford, E., Chang, J.L., Cuomo, C.A., Eichler, E., Fitzgerald, M.G., Jaffe, D.B., et al. 2006. Human chromosome 11 DNA sequence and analysis including novel gene identification. *Nature* 440: 497-500.
5. Allen, D.G. and Whitehead, N.P. 2010. Duchenne muscular dystrophy—what causes the increased membrane permeability in skeletal muscle? *Int. J. Biochem. Cell Biol.* 43: 290-294.
6. Jeon, H., Go, Y., Seo, M., Lee, W.H. and Suk, K. 2010. Functional selection of phagocytosis-promoting genes: cell sorting-based selection. *J. Biomol. Screen.* 15: 949-955.
7. Tsakadze, N., Katzin, L.W., Krishnan, S. and Behrouz, R. 2010. Cerebral infarction in Duchenne muscular dystrophy. *J. Stroke Cerebrovasc. Dis.* 20: 264-265.

CHROMOSOMAL LOCATION

Genetic locus: Pamr1 (mouse) mapping to 2 E2.

PRODUCT

PAMR1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PAMR1 shRNA Plasmid (m): sc-143257-SH and PAMR1 shRNA (m) Lentiviral Particles: sc-143257-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PAMR1 siRNA (m) is recommended for the inhibition of PAMR1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PAMR1 gene expression knockdown using RT-PCR Primer: PAMR1 (m)-PR: sc-143257-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.