



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



epsin 4 siRNA (m): sc-144916

BACKGROUND

The mechanism by which receptor tyrosine kinases (RTKs) modulate cellular physiology in response to stimuli is critical to the understanding of growth regulation. Errors in RTK signaling pathways may result in cellular transformation and, ultimately, in cancer. Two novel EGF receptor substrates function in this pathway, designated EGF-receptor pathway substrates 8 and 15, or Eps8 and Eps15. Epsin is a binding partner to Eps15. Both epsin and Eps15 have a ubiquitous tissue distribution but are concentrated in presynaptic nerve terminals specialized for the Clathrin-mediated endocytosis of synaptic vesicles. Disruption of epsin function blocks Clathrin-mediated endocytosis. Epsin, along with its binding partner Eps15, is proposed to be involved in the assistance of Clathrin coat rearrangement during Clathrin coated pit invagination. The Epsin 4 gene is located on chromosome 5q33, and encodes for a clathrin-associated member of the epsin family that has a role in transport and stability of neurotransmitter vesicles at the synapses and within neurons. Abnormalities in the structure, function, or expression of epsin 4 are linked to schizophrenia susceptibility.

REFERENCES

1. Ford, M.G., Mills, I.G., Peter, B.J., Vallis, Y., Praefcke, G.J., Evans, P.R. and McMahon, H.T. 2002. Curvature of clathrin-coated pits driven by epsin. *Nature* 419: 361-366.
2. Stahelin, R.V., Long, F., Peter, B.J., Murray, D., De Camilli, P., McMahon, H.T. and Cho, W. 2003. Contrasting membrane interaction mechanisms of AP180 N-terminal homology (ANTH) and epsin N-terminal homology (ENTH) domains. *J. Biol. Chem.* 278: 28993-28999.
3. Brzustowicz, L.M., Simone, J., Mohseni, P., Hayter, J.E., Hodgkinson, K.A., Chow, E.W. and Bassett, A.S. 2004. Linkage disequilibrium mapping of schizophrenia suscept region of chromosome 1q22. *Am. J. Hum. Genet.* 74: 1057-1063.
4. Hyun, T.S., Rao, D.S., Saint-Dic, D., Michael, L.E., Kumar, P.D., Bradley, S.V., Mizukami, I.F., Oravec-Wilson, K.I. and Ross, T.S. 2004. HIP1 and HIP1r stabilize recep via epsin N-terminal homology domains. *J. Biol. Chem.* 279: 14294-14306.
5. Pimm, J., McQuillin, A., Thirumalai, S., Lawrence, J., Quested, D., Bass, N., Lamb, G., Moorey, H., Datta, S.R., Kalsi, G., Badacsonyi, A., Kelly, K., Morgan, J., Punukollu, B., Curtis, D. and Gurling, H. 2005. The epsin 4 gene on chromosome 5 enthoproin, is involved in the genetic susceptibility to schizophrenia. *Am. J. Hum. Genet.* 76: 902-907.
6. Kweon, D.H., Shin, Y.K., Shin, J.Y., Lee, J.H., Lee, J.B., Seo, J.H. and Kim, Y.S. 2006. Membrane topology of helix 0 of the Epsin N-terminal homology domain. *Mol. Cells* 21: 428-435.
7. Liou, Y.J., Lai, I.C., Wang, Y.C., Bai, Y.M., Lin, C.C., Lin, C.Y., Chen, T.T. and Chen, J.Y. 2006. Genetic analysis of the human ENTH (epsin 4) gene and schizophrenia. *Schizophr. Res.* 84: 236-243.
8. Tang, R.Q., Zhao, X.Z., Shi, Y.Y., Tang, W., Gu, N.F., Feng, G.Y., Xing, Y.L., Zhu, S.M., Sang, H., Liang, P.J. and He, L. 2006. Family-based association study of epsin 4 and schizophrenia. *Mol. Psychiatry* 11: 395-399.

CHROMOSOMAL LOCATION

Genetic locus: Clint1 (mouse) mapping to 11 B1.1.

PRODUCT

epsin 4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see epsin 4 shRNA Plasmid (m): sc-144916-SH and epsin 4 shRNA (m) Lentiviral Particles: sc-144916-V as alternate gene silencing products.

For independent verification of epsin 4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-144916A, sc-144916B and sc-144916C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

epsin 4 siRNA (m) is recommended for the inhibition of epsin 4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor epsin 4 gene expression knockdown using RT-PCR Primer: epsin 4 (m)-PR: sc-144916-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.