



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

eRF3b siRNA (m): sc-144925

BACKGROUND

eRF3b (eukaryotic peptide chain release factor subunit 3b), also known as Gsp2 (G₁ to S phase transition 2), is a 632 amino acid cytoplasmic protein belonging to the GTP-binding elongation factor family. Highly expressed in brain, eRF3b is involved in translation termination in response to the termination codons UAA, UAG and UGA. As a potent stimulator, eRF3b may play a role in the release factor activity of eRF1. eRF3b exhibits GTPase activity, which is ribosome- and eRF1-dependent, and may participate in cell cycle progression. eRF3b is moderately expressed in spleen and lung with weak expression in heart, liver and kidney. eRF3b contains three GTP binding sites and is a component of the mRNA surveillance SURF complex.

REFERENCES

1. Wallrapp, C., et al. 1997. Characterization of a high copy number amplification at 6q24 in pancreatic cancer identifies c-Myb as a candidate oncogene. *Cancer Res.* 57: 3135-3139.
2. Hoshino, S., et al. 1998. Molecular cloning of a novel member of the eukaryotic polypeptide chain-releasing factors (eRF). Its identification as eRF3 interacting with eRF1. *J. Biol. Chem.* 273: 22254-22259.
3. Jakobsen, C.G., et al. 2001. Identification of a novel termination release factor eRF3b expressing the eRF3 activity *in vitro* and *in vivo*. *Mol. Biol.* 35: 672-681.
4. Le Goff, C., et al. 2002. Mouse GSPT2, but not GSPT1, can substitute for yeast eRF3 *in vivo*. *Genes Cells* 7: 1043-1057.
5. Chauvin, C., et al. 2005. Involvement of human release factors eRF3a and eRF3b in translation termination and regulation of the termination complex formation. *Mol. Cell. Biol.* 25: 5801-5811.
6. Zhouravleva, G., et al. 2006. Evolution of translation termination factor eRF3: is GSPT2 generated by retrotransposition of GSPT1's mRNA? *IUBMB Life* 58: 199-202.
7. Zhuravleva, G.A., et al. 2007. Conservation of the MC domains in eukaryotic termination factor eRF3. *Genetika* 43: 38-44.
8. Chauvin, C., et al. 2007. Human eukaryotic release factor 3a depletion causes cell cycle arrest at G₁ phase through inhibition of the mTOR pathway. *Mol. Cell. Biol.* 27: 5619-5629.
9. Tarasov, O.V., et al. 2008. Evaluation of the gene encoding translation termination factor eRF3 as a possible phylogenetic marker. *Mol. Biol.* 42: 937-946.

CHROMOSOMAL LOCATION

Genetic locus: Gsp2 (mouse) mapping to X C3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

eRF3b siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see eRF3b shRNA Plasmid (m): sc-144925-SH and eRF3b shRNA (m) Lentiviral Particles: sc-144925-V as alternate gene silencing products.

For independent verification of eRF3b (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-144925A, sc-144925B and sc-144925C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

eRF3b siRNA (m) is recommended for the inhibition of eRF3b expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

eRF3b (32): sc-135899 is recommended as a control antibody for monitoring of eRF3b gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor eRF3b gene expression knockdown using RT-PCR Primer: eRF3b (m)-PR: sc-144925-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.