



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



RMDN3 siRNA (m): sc-145057

BACKGROUND

RMDN3 (regulator of microtubule dynamics protein 3), also known as FAM82A2 (family with sequence similarity 82, member A2), FAM82C, PTPIP51 (protein tyrosine phosphatase-interacting protein 51) or cerebral protein 10, is a 470 amino acid protein belonging to the FAM82/RMD family that may play a role in differentiation and apoptosis of keratinocytes. In addition to exhibiting nuclear and mitochondrial membrane localization, RMDN3 also localizes to cytoplasm during interphase and spindle microtubules and spindle poles during mitosis. While widely expressed, RMDN3 is found at high levels in epidermis, seminiferous epithelium and fast twitch fibers of skeletal muscle. RMDN3 has also been found in hippocampal neurons, axons of the peripheral nervous system and ganglion cells of the autonomic nervous system. RMDN3 may function as a cellular signaling partner in angiogenesis and vascular remodeling, and overexpression of RMDN3 is known to induce apoptosis.

REFERENCES

1. Stenzinger, A., et al. 2005. The novel protein PTPIP51 exhibits tissue- and cell-specific expression. *Histochem. Cell Biol.* 123: 19-28.
2. Lv, B.F., et al. 2006. Protein tyrosine phosphatase interacting protein 51 (PTPIP51) is a novel mitochondria protein with an N-terminal mitochondrial targeting sequence and induces apoptosis. *Apoptosis* 11: 1489-1501.
3. Oishi, K., et al. 2007. RMD-1, a novel microtubule-associated protein, functions in chromosome segregation in *Caenorhabditis elegans*. *J. Cell Biol.* 179: 1149-1162.
4. Koch, P., et al. 2008. The novel protein PTPIP51 is expressed in human keratinocyte carcinomas and their surrounding stroma. *J. Cell. Mol. Med.* 12: 2083-2095.

CHROMOSOMAL LOCATION

Genetic locus: *Rmdn3* (mouse) mapping to 2 E5.

PRODUCT

RMDN3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RMDN3 shRNA Plasmid (m): sc-145057-SH and RMDN3 shRNA (m) Lentiviral Particles: sc-145057-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RMDN3 siRNA (m) is recommended for the inhibition of RMDN3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RMDN3 gene expression knockdown using RT-PCR Primer: RMDN3 (m)-PR: sc-145057-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.