



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



FANK1 siRNA (m): sc-145066

BACKGROUND

FANK1 (Fibronectin type III and ankyrin repeat domains 1), also known as HSD13, is a 345 amino acid nuclear and cytoplasmic testis-specific protein found primarily in pachytene spermatocytes and round spermatids. Containing six ANK repeats and a single Fibronectin type-III domain, FANK1 undergoes alternative splicing events to form three isoforms. Possessing DNA binding activity, FANK1 is suggested to act as a transcription factor and may regulate gene expression during spermatogenesis. The gene encoding FANK1 maps to human chromosome 10, which contains over 800 genes and 135 million nucleotides, making up nearly 4.5% of the human genome. PTEN is an important tumor suppressor gene located on chromosome 10 and, when defective, causes a genetic predisposition to cancer development known as Cowden syndrome.

REFERENCES

1. Teresi, R.E., Zbuk, K.M., Pezzolesi, M.G., Waite, K.A. and Eng, C. 2007. Cowden syndrome-affected patients with PTEN promoter mutations demonstrate abnormal protein translation. *Am. J. Hum. Genet.* 81: 756-767.
2. Zheng, Z., Zheng, H. and Yan, W. 2007. FANK1 is a testis-specific gene encoding a nuclear protein exclusively expressed during the transition from the meiotic to the haploid phase of spermatogenesis. *Gene Expr. Patterns* 7: 777-783.
3. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611640. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Tremblay, K., Lemire, M., Potvin, C., Tremblay, A., Hunninghake, G.M., Raby, B.A., Hudson, T.J., Perez-Iratxeta, C., Andrade-Navarro, M.A. and Laprise, C. 2008. Genes to diseases (G2D) computational method to identify asthma candidate genes. *PLoS ONE* 3: e2907.
5. Trynka, G., Zhernakova, A., Romanos, J., Franke, L., Hunt, K.A., Turner, G., Bruinenberg, M., Heap, G.A., Platteel, M., Ryan, A.W., de Kovel, C., Holmes, G.K., Howdle, P.D., Walters, J.R., Sanders, D.S., Mulder, C.J., et al. 2009. Coeliac disease-associated risk variants in TNFAIP3 and REL implicate altered NFκB signalling. *Gut* 58: 1078-1083.

CHROMOSOMAL LOCATION

Genetic locus: Fank1 (mouse) mapping to 7 F3.

PRODUCT

FANK1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FANK1 shRNA Plasmid (m): sc-145066-SH and FANK1 shRNA (m) Lentiviral Particles: sc-145066-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FANK1 siRNA (m) is recommended for the inhibition of FANK1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

FANK1 (F-5): sc-398026 is recommended as a control antibody for monitoring of FANK1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FANK1 gene expression knockdown using RT-PCR Primer: FANK1 (m)-PR: sc-145066-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.