



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# FAR2 siRNA (m): sc-145071

## BACKGROUND

The conversion of fatty acids to fatty alcohols is required for the synthesis of wax monoesters and ether lipids. Fatty acyl-CoA reductase 2 (FAR2), also known as male sterility domain-containing protein 1 (MLSTD1), is a 515 amino acid protein that catalyzes the reduction of saturated fatty acyl-CoA with chain length C16, C18 and C10-C14 to fatty alcohols. FAR2, which is a member of the fatty acyl-CoA reductase family, is a multi-pass membrane protein and has been shown to localize to the peroxisome and the endoplasmic reticulum. The highest levels of FAR2 were detected in brain and eyelid, which contains wax-laden meibomian glands. The gene encoding FAR2 is located on chromosome 12, which comprises nearly 4.5% of the human genome.

## REFERENCES

1. Wang, X., et al. 1995. Solubilization, purification and characterization of fatty acyl-CoA reductase from duck uropygial gland. *Biochem. Biophys. Res. Commun.* 208: 210-215.
2. Hajra, A.K., et al. 1996. Lipid biosynthesis in peroxisomes. *Ann. N.Y. Acad. Sci.* 804: 129-141.
3. Vioque, J., et al. 1997. Resolution and purification of an aldehyde-generating and an alcohol-generating fatty acyl-CoA reductase from pea leaves (*Pisum sativum* L.). *Arch. Biochem. Biophys.* 340: 64-72.
4. Yamashita, A., et al. 1997. Acyltransferases and transacylases involved in fatty acid remodeling of phospholipids and metabolism of bioactive lipids in mammalian cells. *J. Biochem.* 122: 1-16.
5. Phipps, A.N., et al. 2000. Peroxisome distribution along the crypt-villus axis of the guinea pig small intestine. *Mol. Cell. Biochem.* 203: 119-126.
6. Cheng, J.B., et al. 2004. Mammalian wax biosynthesis. I. Identification of two fatty acyl-Coenzyme A reductases with different substrate specificities and tissue distributions. *J. Biol. Chem.* 279: 37789-37797.
7. Cheng, J.B. and Russell, D.W. 2004. Mammalian wax biosynthesis. II. Expression cloning of wax synthase cDNAs encoding a member of the acyltransferase enzyme family. *J. Biol. Chem.* 279: 37798-37807.
8. Costaglioli, P., et al. 2005. Profiling candidate genes involved in wax biosynthesis in *Arabidopsis thaliana* by microarray analysis. *Biochim. Biophys. Acta* 1734: 247-258.
9. Doan, T.T., et al. 2009. Functional expression of five *Arabidopsis* fatty acyl-CoA reductase genes in *Escherichia coli*. *J. Plant Physiol.* 166: 787-796.

## CHROMOSOMAL LOCATION

Genetic locus: Far2 (mouse) mapping to 6 G3.

## PRODUCT

FAR2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FAR2 shRNA Plasmid (m): sc-145071-SH and FAR2 shRNA (m) Lentiviral Particles: sc-145071-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

FAR2 siRNA (m) is recommended for the inhibition of FAR2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FAR2 gene expression knockdown using RT-PCR Primer: FAR2 (m)-PR: sc-145071-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.