



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FBXO22 siRNA (m): sc-145110

BACKGROUND

FBXO22 (F-box only protein 22), also known as FBX22 or FBX22p44, is a 403 amino acid protein that is predominately expressed in the liver. F-box proteins are critical components of the SCF (Skp1-CUL-1-F-box protein) type E3 ubiquitin ligase complex and are involved in substrate recognition and recruitment for ubiquitination. They are members of a larger family of proteins that are involved in the regulation of a wide variety of cellular processes (including the cell cycle, immune response, signaling cascades and developmental processes) through the targeting of proteins, such as cyclins and cyclin-dependent kinase inhibitors (CDKNs), I κ B- α and β -catenin, for degradation by the proteasome after ubiquitination. Three isoforms of FBXO22 exist due to alternative splicing.

REFERENCES

1. Winston, J.T., Koepp, D.M., Zhu, C., Elledge, S.J. and Harper, J.W. 1999. A family of mammalian F-box proteins. *Curr. Biol.* 9: 1180-1182.
2. Cenciarelli, C., Chiaur, D.S., Guardavaccaro, D., Parks, W., Vidal, M. and Pagano, M. 1999. Identification of a family of human F-box proteins. *Curr. Biol.* 9: 1177-1179.
3. Jin, J., Cardozo, T., Lovering, R.C., Elledge, S.J., Pagano, M. and Harper, J.W. 2004. Systematic analysis and nomenclature of mammalian F-box proteins. *Genes Dev.* 18: 2573-2580.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 609096. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Yoshida, Y. 2007. F-box proteins that contain sugar-binding domains. *Biosci. Biotechnol. Biochem.* 71: 2623-2631.
6. Cooke, P.S., Holsberger, D.R., Cimafranca, M.A., Meling, D.D., Beals, C.M., Nakayama, K., Nakayama, K.I. and Kiyokawa, H. 2007. The F box protein S phase kinase-associated protein 2 regulates adipose mass and adipocyte number *in vivo*. *Obesity* 15: 1400-1408.
7. Bernis, C., Vigneron, S., Burgess, A., Labbe, J.C., Fesquet, D., Castro, A. and Lorca, T. 2007. Pin1 stabilizes Emi1 during G₂ phase by preventing its association with SCF^{FBTRCP}. *EMBO Rep.* 8: 91-98.
8. SWISS-PROT/TrEMBL (Q8NEZ5). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: Fbxo22 (mouse) mapping to 9 B.

PRODUCT

FBXO22 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FBXO22 shRNA Plasmid (m): sc-145110-SH and FBXO22 shRNA (m) Lentiviral Particles: sc-145110-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FBXO22 siRNA (m) is recommended for the inhibition of FBXO22 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FBXO22 gene expression knockdown using RT-PCR Primer: FBXO22 (m)-PR: sc-145110-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.