



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Fibulin-7 siRNA (m): sc-145177

BACKGROUND

Fibulin proteins contribute to normal development of elastic fiber systems in various types of organs that require elasticity, such as vasculature, lung and skin. Fibulin-7, also known as TM14 or FBLN7, is a 439 amino acid extracellular matrix protein that belongs to the Fibulin family. Containing two EGF-like domains and one sushi (CCP/SCR) domain, Fibulin-7 exists as four alternatively spliced isoforms. Fibulin-7 is considered an adhesion protein that interacts with extracellular matrix molecules in developing teeth, and may be involved in differentiation and maintenance of odontoblasts as well as in dentin formation. Fibulin-7 is post-translationally glycosylated with N-linked oligosaccharides and interacts with heparin, fibronectin, Fibulin-1 and DSP (dentin sialophosphoprotein). Fibulin-7 is encoded by a gene located on human chromosome 2q13, which consists of 237 million bases, encodes over 1,400 genes and makes up approximately 8% of the human genome.

REFERENCES

1. Giltay, R., Timpl, R. and Kostka, G. 1999. Sequence, recombinant expression and tissue localization of two novel extracellular matrix proteins, Fibulin-3 and Fibulin-4. *Matrix Biol.* 18: 469-480.
2. Gallagher, W.M., Greene, L.M., Ryan, M.P., Sierra, V., Berger, A., Laurent-Puig, P. and Conseiller, E. 2001. Human Fibulin-4: analysis of its biosynthetic processing and mRNA expression in normal and tumour tissues. *FEBS Lett.* 489: 59-66.
3. El-Hallous, E., Sasaki, T., Hubmacher, D., Getie, M., Tiedemann, K., Brinckmann, J., Bätge, B., Davis, E.C. and Reinhardt, D.P. 2007. Fibrillin-1 interactions with Fbulins depend on the first hybrid domain and provide an adaptor function to tropoelastin. *J. Biol. Chem.* 282: 8935-8946.
4. de Vega, S., Iwamoto, T., Nakamura, T., Hozumi, K., McKnight, D.A., Fisher, L.W., Fukumoto, S. and Yamada, Y. 2007. TM14 is a new member of the Fibulin family (Fibulin-7) that interacts with extracellular matrix molecules and is active for cell binding. *J. Biol. Chem.* 282: 30878-30888.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611551. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Arany, S., Koyota, S. and Sugiyama, T. 2009. Nerve growth factor promotes differentiation of odontoblast-like cells. *J. Cell. Biochem.* 106: 539-545.
7. Lisi, S., D'Amore, M., Scagliusi, P., Mitolo, V. and Sisto, M. 2009. Anti-Ro/SSA autoantibody-mediated regulation of extracellular matrix Fbulins in human epithelial cells of the salivary gland. *Scand. J. Rheumatol.* 38: 198-206.

CHROMOSOMAL LOCATION

Genetic locus: Fbln7 (mouse) mapping to 2 F1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Fibulin-7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Fibulin-7 shRNA Plasmid (m): sc-145177-SH and Fibulin-7 shRNA (m) Lentiviral Particles: sc-145177-V as alternate gene silencing products.

For independent verification of Fibulin-7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145177A, sc-145177B and sc-145177C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Fibulin-7 siRNA (m) is recommended for the inhibition of Fibulin-7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Fibulin-7 gene expression knockdown using RT-PCR Primer: Fibulin-7 (m)-PR: sc-145177-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.