



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FNDC4 siRNA (m): sc-145213

BACKGROUND

Fibronectins are multi-domain glycoproteins that bind to a variety of substances including collagen, Actin, heparin, DNA, fibrin and fibronectin receptors. They are involved in a diverse array of important functions such as blood coagulation, wound healing, cell adhesion, cell differentiation and migration. FNDC4 (fibronectin type III domain-containing protein 4), also known as FRCP1 (fibronectin type III repeat-containing protein 1), is a 234 amino acid membrane protein that contains one fibronectin type-III domain, which serves as a binding site for DNA, heparin or the cell surface. The gene encoding FNDC4 is localized to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome.

REFERENCES

1. Kornblihtt, A.R., Umezawa, K., Vibe-Pedersen, K. and Baralle, F.E. 1985. Primary structure of human fibronectin: differential splicing may generate at least 10 polypeptides from a single gene. *EMBO J.* 4: 1755-1759.
2. Dean, D.C., Bowlus, C.L. and Bourgeois, S. 1987. Cloning and analysis of the promotor region of the human fibronectin gene. *Proc. Natl. Acad. Sci. USA* 84: 1876-1880.
3. Leahy, D.J., Hendrickson, W.A., Aukhil, I. and Erickson, H.P. 1992. Structure of a fibronectin type III domain from tenascin phased by MAD analysis of the selenomethionyl protein. *Science* 258: 987-991.
4. Little, E., Bork, P. and Doolittle, R.F. 1994. Tracing the spread of fibronectin type III domains in bacterial glycohydrolases. *J. Mol. Evol.* 39: 631-643.
5. Teufel, A., Malik, N., Mukhopadhyay, M. and Westphal, H. 2002. Frp1 and Frp2, two novel fibronectin type III repeat containing genes. *Gene* 297: 79-83.
6. Carafoli, F., Saffell, J.L. and Hohenester, E. 2008. Structure of the tandem fibronectin type 3 domains of neural cell adhesion molecule. *J. Mol. Biol.* 377: 524-534.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 611905. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Fndc4 (mouse) mapping to 5 B1.

PRODUCT

FNDC4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FNDC4 shRNA Plasmid (m): sc-145213-SH and FNDC4 shRNA (m) Lentiviral Particles: sc-145213-V as alternate gene silencing products.

For independent verification of FNDC4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145213A, sc-145213B and sc-145213C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FNDC4 siRNA (m) is recommended for the inhibition of FNDC4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FNDC4 gene expression knockdown using RT-PCR Primer: FNDC4 (m)-PR: sc-145213-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.