



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FOXE1 siRNA (m): sc-145224

BACKGROUND

Forkhead box protein E1 (FOXE1) is a member of the forkhead/ winged-helix domain transcription factor family. FOXE1, also designated FKHL15 or TTF-2, complexes with TTF-1 and Pax-8 to induce thyroid follicular cell differentiation and thyroid hormone biosynthesis by regulating the expression of the sodium iodide symporter (NIS), thyroid peroxidase (TPO), thyroglobulin (TG) and the thyrotropin receptor (TSHR). FOXE1 encodes a protein that is expressed in several tissues, including thymus, adult brain, lung, liver, heart and pancreas. The chromosomal location of the FOXE1 gene on 9q22.33 suggests that it may be involved in squamous cell epithelioma and hereditary sensory neuropathy type I. Mutations in the FOXE1 gene lead to the development of congenital hypothyroidism, which occurs in approximately one in four thousand newborns and results in complete or partial failure of thyroid gland development. Patients who are homozygous for a missense mutation in the forkhead domain of the FOXE1 gene can also develop thyroid agenesis, cleft palate and choanal atresia. Subsequently, the FOXE1 gene may be used as a marker to study these disorders.

REFERENCES

1. Chadwick, B.P., et al. 1997. FKHL15, a new human member of the forkhead gene family located on chromosome 9q22. *Genomics* 41: 390-396.
2. Clifton-Bligh, R.J., et al. 1998. Mutation of the gene encoding human TTF-2 associated with thyroid agenesis, cleft palate and choanal atresia. *Nat. Genet.* 19: 399-401.
3. Suzuki, K., et al. 1999. Thyroglobulin regulates follicular function and heterogeneity by suppressing thyroid-specific gene expression. *Biochimie* 81: 329-340.
4. Miyazaki, A., et al. 1999. Tumor necrosis factor α and interferon- γ suppress both gene expression and deoxyribonucleic acid-binding of TTF-2 in FRTL-5 cells. *Endocrinology* 140: 4214-4220.

CHROMOSOMAL LOCATION

Genetic locus: Foxe1 (mouse) mapping to 4 B1.

PRODUCT

FOXE1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FOXE1 shRNA Plasmid (m): sc-145224-SH and FOXE1 shRNA (m) Lentiviral Particles: sc-145224-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FOXE1 siRNA (m) is recommended for the inhibition of FOXE1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contains a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FOXE1 gene expression knockdown using RT-PCR Primer: FOXE1 (m)-PR: sc-145224-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.