



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

G_γ 12 siRNA (m): sc-145285

BACKGROUND

Members of the guanine nucleotide-binding protein (G protein) γ family directly regulate various activities of ion channels and enzymes. Eight known human G protein γ subunits exist, three of which are novel forms that are designated G_γ 4, G_γ 10 and G_γ 11. G_γ 12 (guanine nucleotide binding protein (G protein), γ 12), also known as GNG12, is a 72 amino acid lipid-anchored, cell membrane protein belonging to the G protein γ family. G_γ 12 is essential for GTPase activity, G protein-effector interaction and replacement of GDP by GTP. G_γ 12 may function as a negative regulator of the LPS response and may also be an important factor in the overall inflammatory signaling cascade, which has a central role in many neurodegenerative diseases such as Alzheimer's, AIDS dementia and Parkinson's.

REFERENCES

1. Cali, J.J., Balcueva, E.A., Rybalkin, I. and Robishaw, J.D. 1992. Selective tissue distribution of G protein γ subunits, including a new form of the γ subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
2. Ray, K., Kunsch, C., Bonner, L.M. and Robishaw, J.D. 1995. Isolation of cDNA clones encoding eight different human G protein γ subunits, including three novel forms designated the γ 4, γ 10, and γ 11 subunits. *J. Biol. Chem.* 270: 21765-21771.
3. Downes, G.B. and Gautam, N. 1999. The G protein subunit gene families. *Genomics* 62: 544-552.
4. Hurowitz, E.H., Melnyk, J.M., Chen, Y.J., Kouros-Mehr, H., Simon, M.I. and Shizuya, H. 2000. Genomic characterization of the human heterotrimeric G protein α , β , and γ subunit genes. *DNA Res.* 7: 111-120.
5. Cook, L.A., Schey, K.L., Cleator, J.H., Wilcox, M.D., Dingus, J. and Hildebrandt, J.D. 2001. Identification of a region in G protein γ subunits conserved across species but hypervariable among subunit isoforms. *Protein Sci.* 10: 2548-2555.
6. Liu, P., Zhang, S., Yao, Q., Liu, X., Wang, X., Huang, C., Huang, X., Wang, P., Yuan, M., Liu, J.Y., Wang, Q.K. and Liu, M. 2008. Identification of a genetic locus for autosomal dominant disseminated superficial actinic porokeratosis on chromosome 1p31.3-p31.1. *Hum. Genet.* 123: 507-513.
7. Larson, K.C., Lipko, M., Dabrowski, M. and Draper, M.P. 2010. Gng12 is a novel negative regulator of LPS-induced inflammation in the microglial cell line BV-2. *Inflamm. Res.* 59: 15-22.

CHROMOSOMAL LOCATION

Genetic locus: Gng12 (mouse) mapping to 6 C1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

G_γ 12 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G_γ 12 shRNA Plasmid (m): sc-145285-SH and G_γ 12 shRNA (m) Lentiviral Particles: sc-145285-V as alternate gene silencing products.

For independent verification of G_γ 12 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145285A, sc-145285B and sc-145285C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

G_γ 12 siRNA (m) is recommended for the inhibition of G_γ 12 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G_γ 12 gene expression knockdown using RT-PCR Primer: G_γ 12 (m)-PR: sc-145285-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.