



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

G9a siRNA (m): sc-145298

BACKGROUND

Distinct modifications of histone tails, such as acetylation, phosphorylation and methylation, regulate nuclear processes, such as control of transcription and mitotic chromosome condensation. Histone methyltransferases (HMTases) are among the different groups of enzymes known to catalyze the covalent modification. G9a, a SET domain-containing protein, is a novel mammalian lysine-preferring HMTase. G9a, also known as BAT8, NG36 or HMTase (for mammalian histone methyltransferase), has strong HMTase activity towards Histone H3 Lysine 9 methylation *in vitro*. G9a plays a dominant role in euchromatic Histone H3 Lysine 9 methylation, is essential for early embryogenesis and is involved in the transcriptional repression of developmental genes. Like SUV39H, G9a transfers methyl groups to the lysine residues of Histone H3, but with a 10-20-fold higher activity than SUV39H1. G9a also adds methyl groups to Lysine 27 as well as Lysine 9 in Histone H3. G9a localizes in the nucleus, indicating that it may contribute to the organization of the higher order chromatin structure of non-centromeric loci. The human G9a gene maps to chromosome 6p21.3.

REFERENCES

1. Spies, T., et al. 1989. Human major histocompatibility complex contains a minimum of 19 genes between the complement cluster and HLA-B. *Proc. Natl. Acad. Sci. USA* 86: 8955-8958.
2. Milner, C.M. and Campbell, R.D. 1993. The G9a gene in the human major histocompatibility complex encodes a novel protein containing ankyrin-like repeats. *Biochem. J.* 290: 811-818.
3. Tachibana, M., et al. 2001. Set domain-containing protein, G9a, is a novel lysine-preferring mammalian histone methyltransferase with hyperactivity and specific selectivity to Lysines 9 and 27 of Histone H3. *J. Biol. Chem.* 276: 25309-25317.
4. Brown, S.E., et al. 2001. Novel NG36/G9a gene products encoded within the human and mouse MHC class III regions. *Mamm. Genome* 12: 916-924.
5. Tachibana, M., et al. 2002. G9a histone methyltransferase plays a dominant role in euchromatic Histone H3 Lysine 9 methylation and is essential for early embryogenesis. *Genes Dev.* 16: 1779-1791.

CHROMOSOMAL LOCATION

Genetic locus: Ehmt2 (mouse) mapping to 17 B1.

PRODUCT

G9a siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G9a shRNA Plasmid (m): sc-145298-SH and G9a shRNA (m) Lentiviral Particles: sc-145298-V as alternate gene silencing products.

For independent verification of G9a (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3 nmol of lyophilized siRNA. These include: sc-145298A, sc-145298B and sc-145298C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

G9a siRNA (m) is recommended for the inhibition of G9a expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

G9a (C-9): sc-515726 is recommended as a control antibody for monitoring of G9a gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G9a gene expression knockdown using RT-PCR Primer: G9a (m)-PR: sc-145298-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.