



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GANC siRNA (m): sc-145324

BACKGROUND

A key enzyme in glycogen degradation and metabolism, GANC (glucosidase, α ; neutral C) is a 914 amino acid protein with α -glucosidase activity that belongs to the glycosyl hydrolase 31 family and hydrolyzes non-reducing, terminal 1,4-linked α -D-glucose residues and releases α -D-glucose. The gene encoding GANC maps to human chromosome 15q15.1, a region associated with susceptibility to non-Insulin-dependent (type 2) diabetes mellitus, a disease characterized by high blood glucose levels. Human chromosome 15 houses over 700 genes and comprises nearly 3% of the human genome. Angelman syndrome, Prader-Willi syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in chromosome 15-localized genes.

REFERENCES

1. Martiniuk, F. and Hirschhorn, R. 1980. Human neutral α -glucosidase C: genetic polymorphism including a "null" allele. *Am. J. Hum. Genet.* 32: 497-507.
2. Martiniuk, F., Hirschhorn, R. and Smith, M. 1980. Assignment of the gene for human neutral α -glucosidase C to chromosome 15. *Cytogenet. Cell Genet.* 27: 168-175.
3. Cox, N.J., Frigge, M., Nicolae, D.L., Concannon, P., Hanis, C.L., Bell, G.I. and Kong, A. 1999. Loci on chromosomes 2 (NIDDM1) and 15 interact to increase susceptibility to diabetes in Mexican Americans. *Nat. Genet.* 21: 213-215.
4. Hirschhorn, R., Huie, M.L. and Kasper, J.S. 2002. Computer assisted cloning of human neutral α -glucosidase C (GANC): a new paralog in the glycosyl hydrolase gene family 31. *Proc. Natl. Acad. Sci. USA* 99: 13642-13646.
5. Kawabata, Y., Hata, S., Ono, Y., Ito, Y., Suzuki, K., Abe, K. and Sorimachi, H. 2003. Newly identified exons encoding novel variants of p94/calpain 3 are expressed ubiquitously and overlap the α -glucosidase C gene. *FEBS Lett.* 555: 623-630.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 104180. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Ferrer-Bolufer, I., Dalmau, J., Quiroga, R., Oltra, S., Orellana, C., Monfort, S., Roselló, M., De La Osa, A. and Martinez, F. 2009. Tyrosinemia type 1 and Angelman syndrome due to paternal uniparental isodisomy 15. *J. Inherit. Metab. Dis.* 32: S349-S353.
8. Wawrzik, M., Unmehopa, U.A., Swaab, D.F., van de Nes, J., Buiting, K. and Horsthemke, B. 2010. The C15orf2 gene in the Prader-Willi syndrome region is subject to genomic imprinting and positive selection. *Neurogenetics* 11: 153-161.

CHROMOSOMAL LOCATION

Genetic locus: Ganc (mouse) mapping to 2 E5.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GANC siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GANC shRNA Plasmid (m): sc-145324-SH and GANC shRNA (m) Lentiviral Particles: sc-145324-V as alternate gene silencing products.

For independent verification of GANC (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145324A, sc-145324B and sc-145324C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GANC siRNA (m) is recommended for the inhibition of GANC expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GANC gene expression knockdown using RT-PCR Primer: GANC (m)-PR: sc-145324-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.