



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Glutamine-rich 2 siRNA (m): sc-145453

BACKGROUND

Glutamine-rich 2 is a 1,663 amino acid protein that contains almost a 400 amino acid glutamine-rich region, which is possibly involved in protein-protein interactions. There are three isoforms of Glutamine-rich 2 that are produced as a result of alternative splicing events. The gene encoding Glutamine-rich 2 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome.

REFERENCES

1. Gilbert, F. 1998. Disease genes and chromosomes: disease maps of the human genome. *Chromosome 17. Genet. Test.* 2: 357-381.
2. Knight, J., Munroe, P.B., Pembroke, J.C. and Caulfield, M.J. 2003. Human chromosome 17 in essential hypertension. *Ann. Hum. Genet.* 67: 193-206.
3. Shashi, V., White, J.R., Pettenati, M.J., Root, S.K. and Bell, W.L. 2003. Ring chromosome 17: phenotype variation by deletion size. *Clin. Genet.* 64: 361-365.
4. Barbouti, A., Stankiewicz, P., Nusbaum, C., Cuomo, C., Cook, A., Höglund, M., Johansson, B., Hagemeijer, A., Park, S.S., Mitelman, F., Lupski, J.R. and Fioretos, T. 2004. The breakpoint region of the most common isochromosome, i(17q), in human neoplasia is characterized by a complex genomic architecture with large, palindromic, low-copy repeats. *Am. J. Hum. Genet.* 74: 1-10.
5. Yamamoto, K., Yakushijin, K., Nishikawa, S., Minagawa, K., Katayama, Y., Shimoyama, M. and Matsui, T. 2008. Imatinib resistance in a novel translocation der(17)t(1;17)(q25;p13) with loss of TP53 but without Bcr/Abl kinase domain mutation in chronic myelogenous leukemia. *Cancer Genet. Cytogenet.* 183: 77-81.
6. McCabe, M.G., Ichimura, K., Pearson, D.M., Liu, L., Clifford, S.C., Ellison, D.W. and Collins, V.P. 2009. Novel mechanisms of gene disruption at the medulloblastoma isodiscentric 17p11 breakpoint. *Genes Chromosomes Cancer* 48: 121-131.
7. Dicker, F., Herholz, H., Schnittger, S., Nakao, A., Patten, N., Wu, L., Kern, W., Haferlach, T. and Haferlach, C. 2009. The detection of TP53 mutations in chronic lymphocytic leukemia independently predicts rapid disease progression and is highly correlated with a complex aberrant karyotype. *Leukemia* 23: 117-124.
8. Seifert, H., Mohr, B., Thiede, C., Oelschlägel, U., Schäkel, U., Illmer, T., Soucek, S., Ehninger, G. and Schaich, M. 2009. The prognostic impact of 17p (p53) deletion in 2272 adults with acute myeloid leukemia. *Leukemia* 23: 656-663.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Qrich2 (mouse) mapping to 11 E2.

PRODUCT

Glutamine-rich 2 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Glutamine-rich 2 shRNA Plasmid (m): sc-145453-SH and Glutamine-rich 2 shRNA (m) Lentiviral Particles: sc-145453-V as alternate gene silencing products.

For independent verification of Glutamine-rich 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145453A and sc-145453B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Glutamine-rich 2 siRNA (m) is recommended for the inhibition of Glutamine-rich 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Glutamine-rich 2 gene expression knockdown using RT-PCR Primer: Glutamine-rich 2 (m)-PR: sc-145453-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.