



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Gm221 siRNA (m): sc-145523

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. As a member of the Krüppel C₂H₂-type zinc-finger protein family, ZNF469 (zinc finger protein 469) is a 469 amino acid nuclear protein that contains five C₂H₂-type zinc fingers. ZNF469 is expressed in cornea, sclera, skin fibroblasts and striated muscle. Deleterious mutations in the gene encoding Zinc finger protein 469, are the cause of Ehlers-Danlos syndrome type 6B (EDS6B), also known as brittle cornea syndrome (BCS) or fragilitas oculi with joint hyperextensibility.

REFERENCES

1. Stein, R., Lazar, M. and Adam, A. 1968. Brittle cornea. A familial trait associated with blue sclera. *Am. J. Ophthalmol.* 66: 67-69.
2. Nagase, T., Nakayama, M., Nakajima, D., Kikuno, R. and Ohara, O. 2001. Prediction of the coding sequences of unidentified human genes. XX. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 8: 85-95.
3. Martin, J., Han, C., Gordon, L.A., Terry, A., Prabhakar, S., She, X., Xie, G., Hellsten, U., Chan, Y.M., Altherr, M., Couronne, O., Aerts, A., Bajorek, E., Black, S., Blumer, H., Branscomb, E., Brown, N.C., Bruno, W.J., et al. 2004. The sequence and analysis of duplication-rich human chromosome 16. *Nature* 432: 988-994.
4. Abu, A., Frydman, M., Marek, D., Pras, E., Stolovitch, C., Aviram-Goldring, A., Riesenstein, S., Reznik-Wolf, H. and Pras, E. 2006. Mapping of a gene causing brittle cornea syndrome in Tunisian jews to 16q24. *Invest. Ophthalmol. Vis. Sci.* 47: 5283-5287.
5. Olsen, J.V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P. and Mann, M. 2006. Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127: 635-648.
6. Yu, L.R., Zhu, Z., Chan, K.C., Issaq, H.J., Dimitrov, D.S. and Veenstra, T.D. 2007. Improved titanium dioxide enrichment of phosphopeptides from HeLa cells and high confident phosphopeptide identification by cross-validation of MS/MS and MS/MS/MS spectra. *J. Proteome Res.* 6: 4150-4162.
7. Abu, A., Frydman, M., Marek, D., Pras, E., Nir, U., Reznik-Wolf, H. and Pras, E. 2008. Deleterious mutations in the zinc-finger 469 gene cause brittle cornea syndrome. *Am. J. Hum. Genet.* 82: 1217-1222.

CHROMOSOMAL LOCATION

Genetic locus: Gm221 (mouse) mapping to 10 A1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Gm221 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Gm221 shRNA Plasmid (m): sc-145523-SH and Gm221 shRNA (m) Lentiviral Particles: sc-145523-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Gm221 siRNA (m) is recommended for the inhibition of Gm221 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Gm221 gene expression knockdown using RT-PCR Primer: Gm221 (m)-PR: sc-145523-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.