



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



FAM170A siRNA (m): sc-145634

BACKGROUND

FAM170A (family with sequence similarity 170, member A), also known as ZNFD (zinc finger domain-containing protein), is a 330 amino acid protein containing one C₂H₂-type zinc finger. Existing as four alternatively spliced isoforms, FAM170A is encoded by a gene located on human chromosome 5q23.1. Chromosome 5 consists of 181 million base pairs and comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

1. Rauch, A. and Dörr, H.G. 2007. Chromosome 5q subtelomeric deletion syndrome. *Am. J. Med. Genet. C Semin. Med. Genet.* 145C: 372-376.
2. Villa, N., Redaelli, S., Borroni, C., Colombo, C., Roncaglia, N., Sala, E., Crosti, F., Cappellini, A. and Dalprà, L. 2007. Fetal trisomy 5 mosaicism: case report and literature review. *Am. J. Med. Genet. A* 143A: 2343-2346.
3. Shadduck, R.K., Latsko, J.M., Rossetti, J.M., Haq, B. and Abdulhaq, H. 2007. Recent advances in myelodysplastic syndromes. *Exp. Hematol.* 35: 137-143.
4. Falini, B., Nicoletti, I., Bolli, N., Martelli, M.P., Liso, A., Gorello, P., Mandelli, F., Mecucci, C. and Martelli, M.F. 2007. Translocations and mutations involving the nucleophosmin (NPM1) gene in lymphomas and leukemias. *Haematologica* 92: 519-532.
5. Kristoffersen, K.E. 2008. Speech and language development in cri du chat syndrome: a critical review. *Clin. Linguist. Phon.* 22: 443-457.
6. Valent, P. 2008. Revealing the pathogenesis of the 5q- syndrome. *Eur. J. Clin. Invest.* 38: 539-540.
7. Buysse, K., Crepel, A., Menten, B., Pattyn, F., Antonacci, F., Veltman, J.A., Larsen, L.A., Tümer, Z., de Klein, A., van de Laar, I., Devriendt, K., Mortier, G. and Speleman, F. 2008. Mapping of 5q35 chromosomal rearrangements within a genomically unstable region. *J. Med. Genet.* 45: 672-678.
8. Azman, B.Z., Akhir, S.M., Zilfalil, B.A. and Ankathil, R. 2008. Two cases of deletion 5p syndrome: one with paternal involvement and another with atypical presentation. *Singapore Med. J.* 49: e98-e100.

CHROMOSOMAL LOCATION

Genetic locus: Fam170a (mouse) mapping to 18 D1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

FAM170A siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FAM170A shRNA Plasmid (m): sc-145634-SH and FAM170A shRNA (m) Lentiviral Particles: sc-145634-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FAM170A siRNA (m) is recommended for the inhibition of FAM170A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FAM170A gene expression knockdown using RT-PCR Primer: FAM170A (m)-PR: sc-145634-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.