



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





GMIP siRNA (m): sc-145646

BACKGROUND

The Rho family of GTP-binding proteins plays a role in the development of neuronal structure. The activation of the GTP-bound form is regulated by GTPase-activating proteins, which stimulate GTP hydrolysis, leading to inactivation. GMIP (gem-interacting protein) is a 970 amino acid protein that stimulates the GTPase activity of RhoA *in vitro* and *in vivo*. GMIP interacts with Gem through its N-terminus and has a Rho GTPase-activating protein domain at its C-terminus. GMIP is able to inhibit RhoA function, leading to Actin cytoskeletal reorganization *in vivo*. Encoded by a gene that maps to human chromosome 19p13.11, GMIP contains one phorbol-ester/DAG-type zinc finger and one Rho-GAP domain.

REFERENCES

1. Leone, A., Mitsiades, N., Ward, Y., Spinelli, B., Poulaki, V., Tsokos, M. and Kelly, K. 2001. The Gem GTP-binding protein promotes morphological differentiation in neuroblastoma. *Oncogene* 20: 3217-3225.
2. Aresta, S., de Tand-Heim, M.F., Beranger, F. and de Gunzburg, J. 2002. A novel Rho GTPase-activating-protein interacts with Gem, a member of the Ras superfamily of GTPases. *Biochem. J.* 367: 57-65.
3. Negishi, M. and Katoh, H. 2002. Rho family GTPases as key regulators for neuronal network formation. *J. Biochem.* 132: 157-166.
4. Etienne-Manneville, S. and Hall, A. 2002. Rho GTPases in cell biology. *Nature* 420: 629-635.
5. Tadokoro, K., Hashimoto, R., Tatsumi, M., Kosuga, A., Kamijima, K. and Kunugi, H. 2005. The Gem interacting protein (GMIP) gene is associated with major depressive disorder. *Neurogenetics* 6: 127-133.
6. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609694. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/609694>
7. Hatzoglou, A., Ader, I., Spingard, A., Flanders, J., Saade, E., Leroy, I., Traver, S., Aresta, S. and de Gunzburg, J. 2007. Gem associates with Ezrin and acts via the Rho-GAP protein GMIP to down-regulate the Rho pathway. *Mol. Biol. Cell* 18: 1242-1252.

CHROMOSOMAL LOCATION

Genetic locus: Gmip (mouse) mapping to 8 B3.3.

PRODUCT

GMIP siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GMIP shRNA Plasmid (m): sc-145646-SH and GMIP shRNA (m) Lentiviral Particles: sc-145646-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GMIP siRNA (m) is recommended for the inhibition of GMIP expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GMIP gene expression knockdown using RT-PCR Primer: GMIP (m)-PR: sc-145646-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.