



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GNPTG siRNA (m): sc-145659

BACKGROUND

GNPTG (N-acetylglucosamine-1-phosphate transferase, γ subunit), also known as GlcNAc-1-phosphotransferase subunit γ , UDP-N-acetylglucosamine-1-phosphotransferase subunit γ or C16orf27, is a 305 amino acid secreted protein containing one PRKCSH domain. Localizing to Golgi apparatus, GNPTG is widely expressed and exists as a hexamer of two α , two β and two γ subunits linked by disulfide bonds. The α and β subunits are thought to function in catalytic activity, while the γ subunits may play a role in the recognition of lysosomal enzymes. The hexameric complex making up GNPTG is essential for catalyzing the initial step towards mannose 6-phosphate lysosomal recognition marker synthesis. Mutations to GNPTG have been linked to mucopolipidosis type III complementation group C (MLIIIC), an autosomal recessive disease of lysosomal hydrolase trafficking.

REFERENCES

1. Bao, M., et al. 1996. Bovine UDP-N-acetylglucosamine:lysosomal-enzyme N-acetylglucosamine-1-phosphotransferase. I. Purification and subunit structure. *J. Biol. Chem.* 271: 31437-31445.
2. Bao, M., et al. 1996. Bovine UDP-N-acetylglucosamine:lysosomal-enzyme N-acetylglucosamine-1-phosphotransferase. II. Enzymatic characterization and identification of the catalytic subunit. *J. Biol. Chem.* 271: 31446-31451.
3. Raas-Rothschild, A., et al. 2000. Molecular basis of variant pseudo-hurler polydystrophy (mucopolipidosis IIIC) *J. Clin. Invest.* 105: 673-681.
4. Martin, J., et al. 2004. The sequence and analysis of duplication-rich human chromosome 16. *Nature* 432: 988-994.
5. Pohl, S., et al. 2009. Compensatory expression of human N-acetylglucosaminyl-1-phosphotransferase subunits in mucopolipidosis type III γ . *Biochim. Biophys. Acta* 1792: 221-225.
6. Encarnação, M., et al. 2009. Molecular analysis of the GNPTAB and GNPTG genes in 13 patients with mucopolipidosis type II or type III-identification of eight novel mutations. *Clin. Genet.* 76: 76-84.
7. Persichetti, E., et al. 2009. Identification and molecular characterization of six novel mutations in the UDP-N-acetylglucosamine-1-phosphotransferase γ subunit (GNPTG) gene in patients with mucopolipidosis III γ . *Hum. Mutat.* 30: 978-984.
8. Pohl, S., et al. 2010. Loss of N-acetylglucosamine-1-phosphotransferase γ subunit due to intronic mutation in GNPTG causes mucopolipidosis type III γ : Implications for molecular and cellular diagnostics. *Am. J. Med. Genet. A* 152A: 124-132.
9. Qian, Y., et al. 2010. Functions of the α , β , and γ subunits of UDP-GlcNAc:lysosomal enzyme N-acetylglucosamine-1-phosphotransferase. *J. Biol. Chem.* 285: 3360-3370.

CHROMOSOMAL LOCATION

Genetic locus: Gnptg (mouse) mapping to 17 A3.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GNPTG siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GNPTG shRNA Plasmid (m): sc-145659-SH and GNPTG shRNA (m) Lentiviral Particles: sc-145659-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GNPTG siRNA (m) is recommended for the inhibition of GNPTG expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GNPTG gene expression knockdown using RT-PCR Primer: GNPTG (m)-PR: sc-145659-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.