



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



GnT-I siRNA (m): sc-145661

BACKGROUND

GnT-I (α -1,3-mannosyl-glycoprotein 2- β -N acetylglucosaminyltransferase, GlcNAc-T I, MGAT1) catalyzes the first step in the conversion of oligomannose-type N-glycans to N-acetyl-lactosamine- and hybrid-type N-glycans. GnT-I is a 445 amino acid type II membrane protein that localizes to the Golgi and is essential for normal embryogenesis. GnT-I is expressed ubiquitously and exists as multiple variants that encode the same functional protein.

REFERENCES

1. Kumar, R., Yang, J., Larsen, R.D. and Stanley, P. 1990. Cloning and expression of N-acetylglucosaminyltransferase I, the medial Golgi transferase that initiates complex N-linked carbohydrate formation. *Proc. Natl. Acad. Sci. USA* 87: 9948-9952.
2. Hull, E., Sarkar, M., Spruijt, M.P., Höppener, J.W., Dunn, R. and Schachter, H. 1991. Organization and localization to chromosome 5 of the human UDP-N-acetylglucosamine: α -3-D-mannoside β -1,2-N-acetylglucosaminyltransferase I gene. *Biochem. Biophys. Res. Commun.* 176: 608-615.
3. Kumar, R., Yang, J., Eddy, R.L., Byers, M.G., Shows, T.B. and Stanley, P. 1992. Cloning and expression of the murine gene and chromosomal location of the human gene encoding N-acetylglucosaminyltransferase I. *Glycobiology* 2: 383-393.
4. Tan, J., D'Agostaro, A.F., Bendiak, B., Reck, F., Sarkar, M., Squire, J.A., Leong, P. and Schachter, H. 1995. The human UDP-N-acetylglucosamine: α -6-D-mannoside- β -1,2- N-acetylglucosaminyltransferase II gene (MGAT2). Cloning of genomic DNA, localization to chromosome 14q21, expression in insect cells and purification of the recombinant protein. *Eur. J. Biochem.* 231: 317-328.
5. Yip, B., Chen, S.H., Mulder, H., Höppener, J.W. and Schachter, H. 1997. Organization of the human β -1,2-N-acetylglucosaminyltransferase I gene (MGAT1), which controls complex and hybrid N-glycan synthesis. *Biochem. J.* 321: 465-474.
6. Yen, C.L., Stone, S.J., Cases, S., Zhou, P. and Farese, R.V. 2002. Identification of a gene encoding MGAT1, a monoacylglycerol acyltransferase. *Proc. Natl. Acad. Sci. USA* 99: 8512-8517.

CHROMOSOMAL LOCATION

Genetic locus: Mgat1 (mouse) mapping to 11 B1.2.

PRODUCT

GnT-I siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GnT-I shRNA Plasmid (m): sc-145661-SH and GnT-I shRNA (m) Lentiviral Particles: sc-145661-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GnT-I siRNA (m) is recommended for the inhibition of GnT-I expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GnT-I gene expression knockdown using RT-PCR Primer: GnT-I (m)-PR: sc-145661-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.