



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GOLT1A siRNA (m): sc-145673

BACKGROUND

The Golgi complex plays an essential role in the post-translational modification and sorting of proteins transported from the endoplasmic reticulum (ER). The Golgi stack consists of a distinct cis face, or entry face, and a *trans* face, or exit face, which are connected via the cis, medial and trans Golgi networks. GOLT1A (Golgi transport 1A), also known as hGOT1b or vesicle transport protein GOT1A, is a 132 amino acid multi-pass membrane protein belonging to the GOT1 family. Localizing to Golgi apparatus, GOLT1A may play a role in the fusion of ER-derived transport vesicles to the Golgi complex. The gene encoding GOLT1A maps to human chromosome 1q32.1 and mouse chromosome 1 E4. Human chromosome 1 spans 260 million base pairs, contains over 3,000 genes, comprises nearly 8% of the human genome and houses a large number of disease-associated genes.

REFERENCES

1. Dobbie, Z., Heinemann, K., Bishop, D.T., Müller, H. and Scott, R.J. 1997. Identification of a modifier gene locus on chromosome 1p35-36 in familial adenomatous polyposis. *Hum. Genet.* 99: 653-657.
2. Eudy, J.D., Weston, M.D., Yao, S., Hoover, D.M., Rehm, H.L., Ma-Edmonds, M., Yan, D., Ahmad, I., Cheng, J.J., Ayuso, C., Cremers, C., Davenport, S., Moller, C., Talmadge, C.B., Beisel, K.W., Tamayo, M., et al. 1998. Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type IIa. *Science* 280: 1753-1757.
3. Conchon, S., Cao, X., Barlowe, C. and Pelham, H.R. 1999. Got1p and Sft2p: membrane proteins involved in traffic to the Golgi complex. *EMBO J.* 18: 3934-3946.
4. Ríos, R.M., Sanchís, A., Tassin, A.M., Fedriani, C. and Bornens, M. 2004. GMAP-210 recruits γ -Tubulin complexes to *cis*-Golgi membranes and is required for Golgi ribbon formation. *Cell* 118: 323-335.
5. Pasilova, M., Russell, A.M., Wanner, A., Wolf, A., Dobbie, Z., Müller, H.J. and Heinemann, K. 2004. Exclusion of an extracolonic disease modifier locus on chromosome 1p33-36 in a large Swiss familial adenomatous polyposis kindred. *Eur. J. Hum. Genet.* 12: 365-371.
6. Oliveira, S.A., Li, Y.J., Nouredine, M.A., Zuchner, S., Qin, X., Pericak-Vance, M.A. and Vance, J.M. 2005. Identification of risk and age-at-onset genes on chromosome 1p in Parkinson disease. *Am. J. Hum. Genet.* 77: 252-264.
7. Ouchi, M., West, K., Crabb, J.W., Kinoshita, S. and Kamei, M. 2005. Proteomic analysis of vitreous from diabetic macular edema. *Exp. Eye Res.* 81: 176-182.
8. Barr, F.A. and Egerer, J. 2005. Golgi positioning: are we looking at the right MAP? *J. Cell Biol.* 168: 993-998.
9. Yokoi, T., Koide, R., Matsuo, K., Nakagawa, A. and Azuma, N. 2009. Analysis of the vitreous membrane in a case of type 1 Stickler syndrome. *Graefes Arch. Clin. Exp. Ophthalmol.* 247: 715-718.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Golt1a (mouse) mapping to 1 E4.

PRODUCT

GOLT1A siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GOLT1A shRNA Plasmid (m): sc-145673-SH and GOLT1A shRNA (m) Lentiviral Particles: sc-145673-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GOLT1A siRNA (m) is recommended for the inhibition of GOLT1A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GOLT1A gene expression knockdown using RT-PCR Primer: GOLT1A (m)-PR: sc-145673-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.