



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



GPR-137 siRNA (m): sc-145689

BACKGROUND

G protein-coupled receptors (GPRs or GPCRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, are members of the largest protein family and play a role in many different stimulus-response pathways. G protein-coupled receptors mediate extracellular signals into intracellular signals (G protein activation). They respond to a wide variety of signaling molecules, including hormones, neurotransmitters and other proteins and peptides. GPR proteins are usually integral seven pass membrane proteins with some conserved amino acid regions. GPR-137 (G protein-coupled receptor 137), also known as TM7SF1L1 (transmembrane 7 superfamily member 1-like 1 protein), C11orf4 or GPR-137A, is a 417 amino acid multi-pass membrane protein that belongs to the GPR-137 family. Existing as three alternatively spliced isoforms, the gene encoding GPR-137 maps to human chromosome 11q13.1.

REFERENCES

1. O'Brien, K.P., Tapia-Páez, I., Stahle-Bäckdahl, M., Kedra, D. and Dumanski, J.P. 2000. Characterization of five novel human genes in the 11q13-q22 region. *Biochem. Biophys. Res. Commun.* 273: 90-94.
2. Covington, D.K., Briscoe, C.A., Brown, A.J. and Jayawickreme, C.K. 2006. The G protein-coupled receptor 40 family (GPR40-GPR43) and its role in nutrient sensing. *Biochem. Soc. Trans.* 34: 770-773.
3. Zaslavsky, A., Singh, L.S., Tan, H., Ding, H., Liang, Z. and Xu, Y. 2006. Homo- and hetero-dimerization of LPA/S1P receptors, OGR1 and GPR4. *Biochim. Biophys. Acta* 1761: 1200-1212.
4. Jones, P.G., Nawoschik, S.P., Sreekumar, K., Uveges, A.J., Tseng, E., Zhang, L., Johnson, J., He, L., Paulsen, J.E., Bates, B. and Pausch, M.H. 2007. Tissue distribution and functional analyses of the constitutively active orphan G protein coupled receptors, GPR26 and GPR78. *Biochim. Biophys. Acta* 1770: 890-901.
5. Yonezawa, T., Kobayashi, Y. and Obara, Y. 2007. Short-chain fatty acids induce acute phosphorylation of the p38 mitogen-activated protein kinase/heat shock protein 27 pathway via GPR43 in the MCF-7 human breast cancer cell line. *Cell. Signal.* 19: 185-193.
6. Rayasam, G.V., Tulasi, V.K., Davis, J.A. and Bansal, V.S. 2007. Fatty acid receptors as new therapeutic targets for diabetes. *Expert Opin. Ther. Targets* 11: 661-671.

CHROMOSOMAL LOCATION

Genetic locus: Gpr137 (mouse) mapping to 19 A.

PRODUCT

GPR-137 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR-137 shRNA Plasmid (m): sc-145689-SH and GPR-137 shRNA (m) Lentiviral Particles: sc-145689-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR-137 siRNA (m) is recommended for the inhibition of GPR-137 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR-137 gene expression knockdown using RT-PCR Primer: GPR-137 (m)-PR: sc-145689-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.