



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GPR108 siRNA (m): sc-145692

BACKGROUND

G protein-coupled receptors (GPRs or GPCRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, are members of the largest protein family and play a role in many different stimulus-response pathways. G protein-coupled receptors mediate extracellular signals into intracellular signals (G protein activation). They respond to a wide variety of signaling molecules, including hormones, neurotransmitters and other proteins and peptides. GPR proteins are usually integral seven pass membrane proteins with some conserved amino acid regions. GPR108 (G protein-coupled receptor 108), also known as LUSTR2 (lung seven transmembrane receptor 2), is a 543 amino acid multi-pass membrane protein that belongs to the LU7TM family. The gene encoding GPR108 maps to human chromosome 19p13.3 and mouse chromosome 17 D.

REFERENCES

1. Grimwood, J., Gordon, L.A., Olsen, A., Terry, A., Schmutz, J., Lamerdin, J., Hellsten, U., Goodstein, D., Couronne, O., Tran-Gyamfi, M., Aerts, A., Altherr, M., Ashworth, L., Bajorek, E., Black, S., Branscomb, E., et al. 2004. The DNA sequence and biology of human chromosome 19. *Nature* 428: 529-535.
2. Covington, D.K., Briscoe, C.A., Brown, A.J. and Jayawickreme, C.K. 2006. The G protein-coupled receptor 40 family (GPR40-GPR43) and its role in nutrient sensing. *Biochem. Soc. Trans.* 34: 770-773.
3. Zaslavsky, A., Singh, L.S., Tan, H., Ding, H., Liang, Z. and Xu, Y. 2006. Homo- and hetero-dimerization of LPA/S1P receptors, OGR1 and GPR4. *Biochim. Biophys. Acta* 1761: 1200-1212.
4. Yang, M., Mailhot, G., Birnbaum, M.J., MacKay, C.A., Mason-Savas, A. and Odgren, P.R. 2006. Expression of and role for ovarian cancer G protein-coupled receptor 1 (OGR1) during osteoclastogenesis. *J. Biol. Chem.* 281: 23598-23605.
5. Jones, P.G., Nawoschik, S.P., Sreekumar, K., Uveges, A.J., Tseng, E., Zhang, L., Johnson, J., He, L., Paulsen, J.E., Bates, B. and Pausch, M.H. 2007. Tissue distribution and functional analyses of the constitutively active orphan G protein-coupled receptors, GPR26 and GPR78. *Biochim. Biophys. Acta* 1770: 890-901.
6. Yonezawa, T., Kobayashi, Y. and Obara, Y. 2007. Short-chain fatty acids induce acute phosphorylation of the p38 mitogen-activated protein kinase/heat shock protein 27 pathway via GPR43 in the MCF-7 human breast cancer cell line. *Cell. Signal.* 19: 185-193.
7. Edgar, A.J. 2007. Human GPR107 and murine Gpr108 are members of the LUSTR family of proteins found in both plants and animals, having similar topology to G protein-coupled receptors. *DNA Seq.* 18: 235-241.
8. Rayasam, G.V., Tulasi, V.K., Davis, J.A. and Bansal, V.S. 2007. Fatty acid receptors as new therapeutic targets for diabetes. *Expert Opin. Ther. Targets* 11: 661-671.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Gpr108 (mouse) mapping to 17 D.

PRODUCT

GPR108 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR108 shRNA Plasmid (m): sc-145692-SH and GPR108 shRNA (m) Lentiviral Particles: sc-145692-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR108 siRNA (m) is recommended for the inhibition of GPR108 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR108 gene expression knockdown using RT-PCR Primer: GPR108 (m)-PR: sc-145692-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.