



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





Grx2 siRNA (m): sc-145788

BACKGROUND

Glutaredoxins, or thioltransferases, are cytoplasmic proteins included in the glutaredoxin family. Human mitochondrial glutaredoxin (Grx2) is the first iron-sulfur containing protein discovered in the glutaredoxin family of proteins. Instrumental in maintaining the redox environment of the cell, Grx2 maintains redox homeostasis specifically within the mitochondrion by allowing the iron-sulfur cluster to function as a redox target for the activation of Grx2 during conditions of stress, when free radicals are formed. Once oxidative stress induces apoptosis in a cell, Grx2 takes action by reducing low molecular weight proteins and disulfides. Any time that NADPH and Glutathione Reductase reside in the cell, Grx2 modulates glutathione-disulfide oxidoreductase activity. The gene that encodes human Grx2 consists of three introns and four exons, and is 10 kilobases long.

REFERENCES

1. Lundberg, M., Johansson, C., Chandra, J., Enoksson, M., Jacobsson, G., Ljung, J., Johansson, M. and Holmgren, A. 2001. Cloning and expression of a novel human glutaredoxin (Grx2) with mitochondrial and nuclear isoforms. *J. Biol. Chem.* 276: 26269-26275.
2. Gladyshev, V.N., Liu, A., Novoselov, S.V., Krysan, K., Sun, Q.A., Kryukov, V.M., Kryukov, G.V. and Lou, M.F. 2001. Identification and characterization of a new mammalian glutaredoxin (thioltransferase), Grx2. *J. Biol. Chem.* 276: 30374-30380.
3. Chung, W.H., Kim, K.D., Cho, Y.J. and Roe, J.H. 2004. Differential expression and role of two dithiol glutaredoxins Grx1 and Grx2 in *Schizosaccharomyces pombe*. *Biochem. Biophys. Res. Commun.* 321: 922-929.
4. Holmgren, A., Johansson, C., Berndt, C., Lönn, M.E., Hudemann, C. and Lillig, C.H. 2005. Thiol redox control via thioredoxin and glutaredoxin systems. *Biochem. Soc. Trans.* 33: 1375-1377.
5. Lundberg, M., Holmgren, A. and Johansson, M. 2005. Human glutaredoxin 2 affinity tag for recombinant peptide and protein purification. *Protein Expr. Purif.* 45: 37-42.
6. Lillig, C.H., Berndt, C., Vergnolle, O., Lönn, M.E., Hudemann, C., Bill, E. and Holmgren, A. 2005. Characterization of human glutaredoxin 2 as iron-sulfur protein: a possible role as redox sensor. *Proc. Natl. Acad. Sci. USA* 102: 8168-8173.
7. Berndt, C., Hudemann, C., Hanschmann, E.M., Axelsson, R., Holmgren, A. and Lillig, C.H. 2006. How does iron-sulfur cluster coordination regulate the activity of human glutaredoxin 2? *Antioxid. Redox Signal.* 9: 151-157.

CHROMOSOMAL LOCATION

Genetic locus: Glrx2 (mouse) mapping to 1 F.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Grx2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Grx2 shRNA Plasmid (m): sc-145788-SH and Grx2 shRNA (m) Lentiviral Particles: sc-145788-V as alternate gene silencing products.

For independent verification of Grx2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145788A, sc-145788B and sc-145788C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Grx2 siRNA (m) is recommended for the inhibition of Grx2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Grx2 gene expression knockdown using RT-PCR Primer: Grx2 (m)-PR: sc-145788-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.