



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HCCR-1 siRNA (m): sc-145903

BACKGROUND

HCCR-1 (human cervical cancer oncogene 1), also known as HCCR-2 or LETMD1, is a 360 amino acid single-pass membrane protein that is expressed in the kidney, liver, skeletal muscle, heart and brain. It is suggested that HCCR-1 may be involved in tumorigenesis and may function as a negative regulator of the p53 tumor suppressor. Overexpression of HCCR-1 may cause mitochondrial dysfunction that can lead to UVC or staurosporine-induced apoptosis resistance and progression of tumor formation. HCCR-1 is considered a candidate biomarker for breast cancer. Various human tumors including leukemia, lymphoma, and carcinomas of the breast, kidney, ovary, stomach, colon, and uterine cervix have high levels of HCCR-1. Six isoforms exist due to alternative splicing events.

REFERENCES

1. Ko, J., Lee, Y.H., Hwang, S.Y., Lee, Y.S., Shin, S.M., Hwang, J.H., Kim, J., Kim, Y.W., Jang, S.W., Ryoo, Z.Y., Kim, I.K., Namkoong, S.E. and Kim, J.W. 2003. Identification and differential expression of novel human cervical cancer oncogene HCCR-2 in human cancers and its involvement in p53 stabilization. *Oncogene* 22: 4679-4689.
2. Ko, J., Shin, S.M., Oh, Y.M., Lee, Y.S., Ryoo, Z.Y., Lee, Y.H., Na, D.S. and Kim, J.W. 2004. Transgenic mouse model for breast cancer: induction of breast cancer in novel oncogene HCCR-2 transgenic mice. *Oncogene* 23: 1950-1953.
3. Jung, S.S., Park, H.S., Lee, I.J., Namkoong, H., Shin, S.M., Cho, G.W., Ha, S.A., Park, Y.G., Lee, Y.S., Ko, J. and Kim, J.W. 2005. The HCCR oncoprotein as a biomarker for human breast cancer. *Clin. Cancer Res.* 11: 7700-7708.
4. Cho, G.W., Shin, S.M., Namkoong, H., Kim, H.K., Ha, S.A., Hur, S.Y., Kim, T.E., Chai, Y.G. and Kim, J.W. 2006. The phosphatidylinositol 3-kinase/Akt pathway regulates the HCCR-1 oncogene expression. *Gene* 384: 18-26.
5. Cho, G.W., Shin, S.M., Kim, H.K., Ha, S.A., Kim, S., Yoon, J.H., Hur, S.Y., Kim, T.E. and Kim, J.W. 2007. HCCR-1, a novel oncogene, encodes a mitochondrial outer membrane protein and suppresses the UVC-induced apoptosis. *BMC Cell Biol.* 8: 50.
6. Yang, Y., Zhang, G.X., Shi, R.H., Lin, Y., Hao, B., Wang, X.Y., Wang, H.D. and Huang, Z.H. 2007. Overexpression of HCCR in hepatocellular carcinoma cells and its clinical significance. *Zhonghua Gan Zang Bing Za Zhi* 15: 223-224.
7. Ha, S.A., Shin, S.M., Lee, Y.J., Kim, S., Kim, H.K., Namkoong, H., Lee, H., Lee, Y.S., Cho, Y.S., Park, Y.G., Jeon, H.M., Oh, C. and Kim, J.W. 2008. HCCRB-1 directly interacting with HCCR-1 induces tumorigenesis through P53 stabilization. *Int. J. Cancer* 122: 501-508.
8. Ha, S.A., Lee, Y.S., Shin, S.M., Kim, H.K., Kim, S., Namkoong, H., Kim, H.J., Jung, S.M., Lee, Y.S., Chung, Y.J., Jung, S.S. and Kim, J.W. 2009. Oncoprotein HCCR-1 expression in breast cancer is well correlated with known breast cancer prognostic factors including the HER2 overexpression, p53 mutation, and ER/PR status. *BMC Cancer* 9: 51.

CHROMOSOMAL LOCATION

Genetic locus: Letmd1 (mouse) mapping to 15 F1.

PRODUCT

HCCR-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HCCR-1 shRNA Plasmid (m): sc-145903-SH and HCCR-1 shRNA (m) Lentiviral Particles: sc-145903-V as alternate gene silencing products.

For independent verification of HCCR-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145903A, sc-145903B and sc-145903C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HCCR-1 siRNA (m) is recommended for the inhibition of HCCR-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HCCR-1 gene expression knockdown using RT-PCR Primer: HCCR-1 (m)-PR: sc-145903-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.