



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





HEG1 siRNA (m): sc-145931

BACKGROUND

HEG1 (HEG homolog 1) is also known as HEG and is a 1,381 amino acid protein that exists as two alternatively spliced isoforms, one of which is a single-pass membrane protein and the other of which is secreted. HEG1, which is expressed in endothelial cells and smooth muscle cells of heart tissue, contains two EGF-like domains that play a role in calcium-binding events which may regulate concentric heart growth. The gene encoding HEG1 maps to human chromosome 3q21.2, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Key tumor suppressing genes on chromosome 3 include those that encode the apoptosis mediator RASSF1, the cell migration regulator HYAL1 and the angiogenesis suppressor SEMA3B. Marfan syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

REFERENCES

1. Mably, J.D., Mohideen, M.A., Burns, C.G., Chen, J.N. and Fishman, M.C. 2003. heart of glass regulates the concentric growth of the heart in zebrafish. *Curr. Biol.* 13: 2138-2147.
2. Bortvin, A., Eggan, K., Skaletsky, H., Akutsu, H., Berry, D.L., Yanagimachi, R., Page, D.C. and Jaenisch, R. 2003. Incomplete reactivation of Oct4-related genes in mouse embryos cloned from somatic nuclei. *Development* 130: 1673-1680.
3. Braga, E.A., Kashuba, V.I., Maliukova, A.V., Loginov, V.I., Senchenko, V.N., Bazov, I.V., Kiselev, L.L. and Zabarovskii, E.R. 2003. New tumor suppressor genes in hot spots of human chromosome 3: new methods of identification. *Mol. Biol.* 37: 194-211.
4. Mably, J.D., Chuang, L.P., Serluca, F.C., Mohideen, M.A., Chen, J.N. and Fishman, M.C. 2006. santa and valentine pattern concentric growth of cardiac myocardium in the zebrafish. *Development* 133: 3139-3146.
5. Picelli, S., Vandrovcova, J., Jones, S., Djureinovic, T., Skoglund, J., Zhou, X.L., Velculescu, V.E., Vogelstein, B. and Lindblom, A. 2008. Genome-wide linkage scan for colorectal cancer susceptibility genes supports linkage to chromosome 3q. *BMC Cancer* 8: 87.
6. Wilting, S.M., de Wilde, J., Meijer, C.J., Berkhof, J., Yi, Y., van Wieringen, W.N., Braakhuis, B.J., Meijer, G.A., Ylstra, B., Snijders, P.J. and Steenbergen, R.D. 2008. Integrated genomic and transcriptional profiling identifies chromosomal loci with altered gene expression in cervical cancer. *Genes Chromosomes Cancer* 47: 890-905.
7. Birch, A.H., Quinn, M.C., Filali-Mouhim, A., Provencher, D.M., Mes-Masson, A.M. and Tonin, P.N. 2008. Transcriptome analysis of serous ovarian cancers identifies differentially expressed chromosome 3 genes. *Mol. Carcinog.* 47: 56-65.
8. Kleaveland, B., Zheng, X., Liu, J.J., Blum, Y., Tung, J.J., Zou, Z., Chen, M., Guo, L., Lu, M.M., Zhou, D., Kitajewski, J., Affolter, M., Ginsberg, M.H. and Kahn, M.L. 2009. Regulation of cardiovascular development and integrity by the heart of glass-cerebral cavernous malformation protein pathway. *Nat. Med.* 15: 169-176.

CHROMOSOMAL LOCATION

Genetic locus: HEG1 (mouse) mapping to 16 B3.

PRODUCT

HEG1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HEG1 shRNA Plasmid (m): sc-145931-SH and HEG1 shRNA (m) Lentiviral Particles: sc-145931-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HEG1 siRNA (m) is recommended for the inhibition of HEG1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HEG1 gene expression knockdown using RT-PCR Primer: HEG1 (m)-PR: sc-145931-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.