



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





Histone cluster 1 H1T siRNA (m): sc-145977

BACKGROUND

Eukaryotic histones are basic and water soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Over 80% of nucleosomes contain the linker Histone H1, derived from an intronless gene, that interacts with linker DNA between nucleosomes and mediates compaction into higher order chromatin. Histones are subject to post-translational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Such modifications include methylation, citrullination, acetylation, phosphorylation, sumoylation, ubiquitination and ADP-ribosylation. Histone cluster 1 H1T, also known as testicular H1 Histone, HIST1H1T or H1TF, is a 207 amino acid protein that contains an H15 (linker Histone H1/H5 globular) domain and belongs to the Histone H1/H5 family. Localizing to the nucleus and chromosome, Histone cluster 1 H1T is a testis specific H1 variant which is expressed during meiosis in spermatogenesis.

REFERENCES

1. Drabent, B., Kardalinos, E. and Doenecke, D. 1991. Structure and expression of the human gene encoding testicular H1 Histone (H1t). *Gene* 103: 263-268.
2. Koppel, D.A., Wolfe, S.A., Fogelfeld, L.A., Merchant, P.S., Prouty, L. and Grimes, S.R. 1994. Primate testicular histone H1t genes are highly conserved and the human H1t gene is located on chromosome 6. *J. Cell. Biochem.* 54: 219-230.
3. Gunjan, A., Paik, J. and Verreault, A. 2005. Regulation of histone synthesis and nucleosome assembly. *Biochimie* 87: 625-635.
4. Rupp, R.A. and Becker, P.B. 2005. Gene regulation by Histone H1: new links to DNA methylation. *Cell* 123: 1178-1179.
5. Bustin, M., Catez, F. and Lim, J.H. 2005. The dynamics of Histone H1 function in chromatin. *Mol. Cell* 17: 617-620.
6. Martin, C. and Zhang, Y. 2005. The diverse functions of histone lysine methylation. *Nat. Rev. Mol. Cell Biol.* 6: 838-849.
7. Wurtele, H. and Verreault, A. 2006. Histone post-translational modifications and the response to DNA double-strand breaks. *Curr. Opin. Cell Biol.* 18: 137-144.
8. Nightingale, K.P., O'Neill, L.P. and Turner, B.M. 2006. Histone modifications: signalling receptors and potential elements of a heritable epigenetic code. *Curr. Opin. Genet. Dev.* 16: 125-136.
9. Hake, S.B. and Allis, C.D. 2006. Histone H3 variants and their potential role in indexing mammalian genomes: the "H3 barcode hypothesis". *Proc. Natl. Acad. Sci. USA* 103: 6428-6435.

CHROMOSOMAL LOCATION

Genetic locus: Hist1h1t (mouse) mapping to 13 A3.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Histone cluster 1 H1T siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Histone cluster 1 H1T shRNA Plasmid (m): sc-145977-SH and Histone cluster 1 H1T shRNA (m) Lentiviral Particles: sc-145977-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Histone cluster 1 H1T siRNA (m) is recommended for the inhibition of Histone cluster 1 H1T expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Histone cluster 1 H1T gene expression knockdown using RT-PCR Primer: Histone cluster 1 H1T (m)-PR: sc-145977-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.