



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Histone cluster 1 H2AC siRNA (m): sc-145980

BACKGROUND

Eukaryotic histones are basic and water soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Two molecules of each of the four core Histones (H2A, H2B, H3 and H4) form the octamer, which is comprised of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. Histones are subject to posttranslational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Histone cluster 2 H2AC (HIST2H2AC), also known as H2A, H2A-GL101 or H2AFQ, is a 129 amino acid member of the Histone H2A family. The gene encoding Histone cluster 2 H2AC is intronless and maps to human chromosome 1q21.2.

REFERENCES

1. Collart, D., Romain, P.L., Huebner, K., Pockwinse, S., Pilapil, S., Cannizzaro, L.A., Lian, J.B., Croce, C.M., Stein, J.L. and Stein, G.S. 1992. A human Histone H2B.1 variant gene, located on chromosome 1, utilizes alternative 3' end processing. *J. Cell. Biochem.* 50: 374-385.
2. Albig, W. and Doenecke, D. 1997. The human histone gene cluster at the D6S105 locus. *Hum. Genet.* 101: 284-294.
3. Marzluff, W.F., Gongidi, P., Woods, K.R., Jin, J. and Maltais, L.J. 2002. The human and mouse replication-dependent histone genes. *Genomics* 80: 487-498.
4. Braastad, C.D., Hovhannisyan, H., van Wijnen, A.J., Stein, J.L. and Stein, G.S. 2004. Functional characterization of a human histone gene cluster duplication. *Gene* 342: 35-40.
5. Wang, H., Wang, L., Erdjument-Bromage, H., Vidal, M., Tempst, P., Jones, R.S. and Zhang, Y. 2004. Role of Histone H2A ubiquitination in Polycomb silencing. *Nature* 431: 873-878.
6. Zhao, Y., Lang, G., Ito, S., Bonnet, J., Metzger, E., Sawatsubashi, S., Suzuki, E., Le Guezennec, X., Stunnenberg, H.G., Krasnov, A., Georgieva, S.G., Schüle, R., Takeyama, K., Kato, S., Tora, L. and Devys, D. 2008. A TIFC/STAGA module mediates Histone H2A and H2B deubiquitination, coactivates nuclear receptors, and counteracts heterochromatin silencing. *Mol. Cell* 29: 92-101.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 602797. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Pinato, S., Scanduzzi, C., Arnaudo, N., Citterio, E., Gaudino, G. and Penengo, L. 2009. RNF168, a new RING finger, MIU-containing protein that modifies chromatin by ubiquitination of Histones H2A and H2AX. *BMC Mol. Biol.* 10: 55.

CHROMOSOMAL LOCATION

Genetic locus: Hist1h2ac (mouse) mapping to 13 A3.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Histone cluster 1 H2AC siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Histone cluster 1 H2AC shRNA Plasmid (m): sc-145980-SH and Histone cluster 1 H2AC shRNA (m) Lentiviral Particles: sc-145980-V as alternate gene silencing products.

For independent verification of Histone cluster 1 H2AC (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-145980A, sc-145980B and sc-145980C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Histone cluster 1 H2AC siRNA (m) is recommended for the inhibition of Histone cluster 1 H2AC expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Histone cluster 1 H2AC gene expression knockdown using RT-PCR Primer: Histone cluster 1 H2AC (m)-PR: sc-145980-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.