



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HMG-4 siRNA (m): sc-146049

BACKGROUND

The HMGB family, whose members include HMG-1, HMG-2, HMG-3 and HMG-4, is a highly conserved group of chromatin-associated proteins. Evidence suggests that the binding of HMG proteins to DNA induces alterations in the DNA architecture including DNA bending and unwinding of the helix. HMG proteins synergize with Oct-2, members of the NFκB family, ATF-2 and c-Jun to activate transcription. Other studies indicate that phosphorylation of HMG proteins is required to stimulate the transcriptional activity of HMG target proteins. HMG proteins bind single-stranded DNA, but are able to induce conformational changes in double-stranded DNA. HMG-4 is a 186 amino acid protein that localizes to the nucleus. Like all other HMGB family proteins, HMG-4 contains two HMG box DNA-binding domains which can bind DNA either in a sequence-specific manner, or without sequence specificity. Additionally, the HMG box DNA-binding domains are able to preferentially bind DNA distortions, such as kinks and bulges, and, via this binding, can bend DNA.

REFERENCES

1. Bustin, M., Lehn, D.A. and Landsman, D. 1990. Structural features of the HMG chromosomal proteins and their genes. *Biochim. Biophys. Acta* 1049: 231-243.
2. Pedersen, T.J., Arwood, L.J., Spiker, S., Guiltinan, M.J. and Thompson, W.F. 1991. High mobility group chromosomal proteins bind to AT-rich tracts flanking plant genes. *Plant Mol. Biol.* 16: 95-104.
3. Putnam, C.D., Copenhaver, G.P., Denton, M.L. and Pikaard, C.S. 1994. The RNA polymerase I transactivator upstream binding factor requires its dimerization domain and high-mobility-group (HMG) box 1 to bend, wrap, and positively supercoil enhancer DNA. *Mol. Cell. Biol.* 14: 6476-6488.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 163905. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Stros, M., Launholt, D. and Grasser, K.D. 2007. The HMG-box: a versatile protein domain occurring in a wide variety of DNA-binding proteins. *Cell. Mol. Life Sci.* 64: 2590-2606.
6. McCauley, M.J., Zimmerman, J., Maher, L.J. and Williams, M.C. 2007. HMGB binding to DNA: single and double box motifs. *J. Mol. Biol.* 374: 993-1004.
7. Watson, M., Stott, K. and Thomas, J.O. 2007. Mapping intramolecular interactions between domains in HMGB1 using a tail-truncation approach. *J. Mol. Biol.* 374: 1286-1297.
8. Kriatchko, A.N., Bergeron, S. and Swanson, P.C. 2008. HMG-box domain stimulation of RAG1/2 cleavage activity is metal ion dependent. *BMC Mol. Biol.* 9: 32.

CHROMOSOMAL LOCATION

Genetic locus: Hmgb4 (mouse) mapping to 4 D2.2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

HMG-4 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HMG-4 shRNA Plasmid (m): sc-146049-SH and HMG-4 shRNA (m) Lentiviral Particles: sc-146049-V as alternate gene silencing products.

For independent verification of HMG-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146049A and sc-146049B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HMG-4 siRNA (m) is recommended for the inhibition of HMG-4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HMG-4 gene expression knockdown using RT-PCR Primer: HMG-4 (m)-PR: sc-146049-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.