



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

hnRNP H2 siRNA (m): sc-146062

BACKGROUND

The Elav-like genes encode for a family of RNA-binding proteins. Elav, a *Drosophila* protein and the first described member, is expressed immediately after neuroblastic differentiation into neurons and is necessary for neuronal differentiation and maintenance. Several mammalian Elav-like proteins, designated HuB (also designated Hel-N1 in human, or Mel-N1 in mouse), HuC and HuD are also expressed in postmitotic neurons. An additional mammalian homolog, HuR, which is also designated HuA, is ubiquitously expressed and is also overexpressed in a wide variety of tumors. Characteristically, these homologs all contain three RNA recognition motifs (RRM) and they specifically bind to AU-rich elements (ARE) in the 3'-untranslated region of mRNAs transcripts. ARE sites target mRNA for rapid degradation and thereby regulate the expression levels of genes involved in cell growth and differentiation. When Elav-like proteins associate with these ARE sites this degradation is inhibited, leading to an increased stability of the corresponding transcript. Elav proteins function within the nucleus, and they are shuttled between the nucleus and cytoplasm by a nuclear export signal, which is a regulatory feature of the Elav-like proteins as it limits their accessibility to ARE sites.

REFERENCES

1. Chagnovich, D., et al. 1996. Differential activity of Elav-like RNA-binding proteins in human neuroblastoma. *J. Biol. Chem.* 271: 33587-33591.
2. Wakamatsu, Y., et al. 1997. Sequential expression and role of Hu RNA-binding proteins during neurogenesis. *Development* 124: 3449-3460.
3. King, P. 1997. Differential expression of the neuroendocrine genes Hel-N1 and HuD in small-cell lung carcinoma: evidence for down-regulation of HuD in the variant phenotype. *Int. J. Cancer* 74: 378-382.
4. Ball, N.S., et al. 1997. Neuron-specific Hel-N1 and HuD as novel molecular markers of neuroblastoma: a correlation of HuD messenger RNA levels with favorable prognostic features. *Clin. Cancer Res.* 3: 1859-1865.
5. Myer, V.E., et al. 1997. Identification of HuR as a protein implicated in AUUUA-mediated mRNA decay. *EMBO J.* 16: 2130-2139.
6. Chung, S., et al. 1997. The Elav-like proteins bind to a conserved regulatory element in the 3'-untranslated region of GAP-43 mRNA. *J. Biol. Chem.* 272: 6593-6598.
7. Peng, S.S., et al. 1998. RNA stabilization by the AU-rich element binding protein, HuR, an Elav protein. *EMBO J.* 17: 3461-3470.
8. Kasashima, K., et al. 1999. Cytoplasmic localization is required for the mammalian Elav-like protein HuD to induce neuronal differentiation. *Genes Cells* 4: 667-683.

CHROMOSOMAL LOCATION

Genetic locus: Hnrnp2 (mouse) mapping to X E3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

hnRNP H2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see hnRNP H2 shRNA Plasmid (m): sc-146062-SH and hnRNP H2 shRNA (m) Lentiviral Particles: sc-146062-V as alternate gene silencing products.

For independent verification of hnRNP H2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146062A, sc-146062B and sc-146062C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

hnRNP H2 siRNA (m) is recommended for the inhibition of hnRNP H2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor hnRNP H2 gene expression knockdown using RT-PCR Primer: hnRNP H2 (m)-PR: sc-146062-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.