



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HSPC121 siRNA (m): sc-146106

BACKGROUND

HSPC121, also known as PTPLAD1 (protein tyrosine phosphatase-like A domain containing 1), BIND1 or HACD3 (3-hydroxyacyl-CoA dehydratase 3), is a 362 amino acid multi-pass membrane protein that belongs to the very long-chain fatty acids dehydratase HACD family. Localizing to the endoplasmic reticulum membrane, HSPC121 is highly expressed in testis, kidney, brain and liver, with lower levels of expression found in skeletal muscle, spleen and heart. HSPC121 contains one CS domain and is induced by AKAP 12, as well as by histone deacetylase inhibitors, and interacts directly Rac 1 and various condensation enzymes of the ELOVL family. Interactions between HSPC121 and the Rac 1 signaling pathway leads to the modulation of gene expression. HSPC121 is also involved in the dehydration step during the synthesis of very long-chain fatty acids (VLCFAs).

REFERENCES

1. Courilleau, D., Chastre, E., Sabbah, M., Redeuilh, G., Atfi, A. and Mester, J. 2000. B-ind1, a novel mediator of Rac1 signaling cloned from sodium butyrate-treated fibroblasts. *J. Biol. Chem.* 275: 17344-17348.
2. Liu, Y., Gao, L. and Gelman, I.H. 2006. SSeCKS/Gravin/AKAP12 attenuates expression of proliferative and angiogenic genes during suppression of v-Src-induced oncogenesis. *BMC Cancer* 6: 105.
3. Sabbah, M., Saucier, C. and Redeuilh, G. 2006. Human B-ind1 gene promoter: cloning and regulation by histone deacetylase inhibitors. *Gene* 374: 128-133.
4. Ikeda, M., Kanao, Y., Yamanaka, M., Sakuraba, H., Mizutani, Y., Igarashi, Y. and Kihara, A. 2008. Characterization of four mammalian 3-hydroxyacyl-CoA dehydratases involved in very long-chain fatty acid synthesis. *FEBS Lett.* 582: 2435-2440.
5. Cantin, G.T., Yi, W., Lu, B., Park, S.K., Xu, T., Lee, J.D. and Yates, J.R. 2008. Combining protein-based IMAC, peptide-based IMAC, and MudPIT for efficient phosphoproteomic analysis. *J. Proteome Res.* 7: 1346-1351.
6. Taguwa, S., Okamoto, T., Abe, T., Mori, Y., Suzuki, T., Moriishi, K. and Matsuura, Y. 2008. Human butyrate-induced transcript 1 interacts with hepatitis C virus NS5A and regulates viral replication. *J. Virol.* 82: 2631-2641.
7. Taguwa, S., Kambara, H., Omori, H., Tani, H., Abe, T., Mori, Y., Suzuki, T., Yoshimori, T., Moriishi, K. and Matsuura, Y. 2009. Cochaperone activity of human butyrate-induced transcript 1 facilitates hepatitis C virus replication through an Hsp90-dependent pathway. *J. Virol.* 83: 10427-10436.
8. Torrado, M., Iglesias, R., Nespereira, B. and Mikhailov, A.T. 2010. Identification of candidate genes potentially relevant to chamber-specific remodeling in postnatal ventricular myocardium. *J. Biomed. Biotechnol.* 2010: 603159.

CHROMOSOMAL LOCATION

Genetic locus: Ptplad1 (mouse) mapping to 9 C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

HSPC121 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HSPC121 shRNA Plasmid (m): sc-146106-SH and HSPC121 shRNA (m) Lentiviral Particles: sc-146106-V as alternate gene silencing products.

For independent verification of HSPC121 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146106A, sc-146106B and sc-146106C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HSPC121 siRNA (m) is recommended for the inhibition of HSPC121 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HSPC121 gene expression knockdown using RT-PCR Primer: HSPC121 (m)-PR: sc-146106-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.