



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HUGT2 siRNA (m): sc-146114

BACKGROUND

HUGT2 is also known as UGCGL2 (UDP-glucose ceramide glucosyltransferase-like 2) or UGT2 (UDP-glucose: glycoprotein glucosyltransferase 2) and is a 1,516 amino acid protein that is expressed abundantly in kidney, pancreas, heart and skeletal muscle. HUGT2 is localized to the lumen of the ER (endoplasmic reticulum) and to the ER-Golgi intermediate compartment which regulates transportation between the ER and the Golgi. HUGT2 is a quality control protein that targets unfolded glycoproteins for glucosylation in a calcium-dependent manner, but may use manganese in the absence of calcium. HUGT2 targets glycans by attaching single glucose residues to glycans with amino acid sequences which are not correctly folded, an enzymatic process that is exhibited during carbohydrate processing. Patients with type 2 diabetes express altered HUGT2 proteins in their β cells, suggesting that these patients may also have defects in their ER glycoprotein quality control abilities.

REFERENCES

1. Arnold, S.M., Fessler, L.I., Fessler, J.H. and Kaufman, R.J. 2000. Two homologues encoding human UDP-glucose: glycoprotein glucosyltransferase differ in mRNA expression and enzymatic activity. *Biochemistry* 39: 2149-2163.
2. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605898. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Arnold, S.M. and Kaufman, R.J. 2003. The noncatalytic portion of human UDP-glucose: glycoprotein glucosyltransferase I confers UDP-glucose binding and transferase function to the catalytic domain. *J. Biol. Chem.* 278: 43320-43328.
4. Marchetti, P., Bugliani, M., Lupi, R., Marselli, L., Masini, M., Boggi, U., Filipponi, F., Weir, G.C., Eizirik, D.L. and Cnop, M. 2007. The endoplasmic reticulum in pancreatic beta cells of type 2 diabetes patients. *Diabetologia* 50: 2486-2494.
5. Mackenzie, P.I., Rogers, A., Treloar, J., Jorgensen, B.R., Miners, J.O. and Meech, R. 2008. Identification of UDP glycosyltransferase 3A1 as a UDP N-acetylglucosaminyltransferase. *J. Biol. Chem.* 283: 36205-36210.

CHROMOSOMAL LOCATION

Genetic locus: Ugg2 (mouse) mapping to 14 E4.

PRODUCT

HUGT2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HUGT2 shRNA Plasmid (m): sc-146114-SH and HUGT2 shRNA (m) Lentiviral Particles: sc-146114-V as alternate gene silencing products.

For independent verification of HUGT2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146114A, sc-146114B and sc-146114C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HUGT2 siRNA (m) is recommended for the inhibition of HUGT2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HUGT2 gene expression knockdown using RT-PCR Primer: HUGT2 (m)-PR: sc-146114-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.