



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



IBtk siRNA (m): sc-146131

BACKGROUND

IBtk (inhibitor of Bruton agammaglobulinemia tyrosine kinase), also known as BTKI or RP1-93K22.1, is a 1,353 amino acid peripheral membrane protein containing three ANK repeats, 2 BTB (POZ) domains and 3 RCC1 repeats. Existing as three alternatively spliced isoforms, IBtk down-regulates BTK kinase activity and disrupts BTK-mediated calcium mobilization, thereby negatively regulating the activation of nuclear factor- κ -B (NF κ B) driven transcription. IBtk isoform 1 is the predominant isoform expressed in all tissues and translocates to the plasma membrane upon IgM stimulation, whereas isoform 2 localizes to the nucleus. Required for B-cell development, IBtk acts as an inhibitor of BTK tyrosine kinase activity. The gene encoding IBtk maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome.

REFERENCES

1. Yamadori, T., Baba, Y., Matsushita, M., Hashimoto, S., Kurosaki, M., Kurosaki, T., Kishimoto, T. and Tsukada, S. 1999. Bruton's tyrosine kinase activity is negatively regulated by Sab, the Btk-SH3 domain-binding protein. *Proc. Natl. Acad. Sci. USA* 96: 6341-6346.
2. Liu, W., Quinto, I., Chen, X., Palmieri, C., Rabin, R.L., Schwartz, O.M., Nelson, D.L. and Scala, G. 2001. Direct inhibition of Bruton's tyrosine kinase by IBtk, a Btk-binding protein. *Nat. Immunol.* 2: 939-946.
3. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606457. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Yu, L., Mohamed, A.J., Simonson, O.E., Vargas, L., Blomberg, K.E., Björkstrand, B., Arteaga, H.J., Nore, B.F. and Smith, C.I. 2008. Proteasome-dependent autoregulation of Bruton tyrosine kinase (Btk) promoter via NF κ B. *Blood* 111: 4617-4626.
5. Spatuzza, C., Schiavone, M., Di Salle, E., Janda, E., Sardiello, M., Fiume, G., Fierro, O., Simonetta, M., Argiriou, N., Faraonio, R., Capparelli, R., Quinto, I. and Scala, G. 2008. Physical and functional characterization of the genetic locus of IBtk, an inhibitor of Bruton's tyrosine kinase: evidence for three protein isoforms of IBtk. *Nucleic Acids Res.* 36: 4402-4416.

CHROMOSOMAL LOCATION

Genetic locus: IBtk (mouse) mapping to 9 E3.1.

PRODUCT

IBtk siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IBtk shRNA Plasmid (m): sc-146131-SH and IBtk shRNA (m) Lentiviral Particles: sc-146131-V as alternate gene silencing products.

For independent verification of IBtk (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146131A, sc-146131B and sc-146131C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IBtk siRNA (m) is recommended for the inhibition of IBtk expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IBtk gene expression knockdown using RT-PCR Primer: IBtk (m)-PR: sc-146131-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.