



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IFN- α 7 siRNA (m): sc-146166

BACKGROUND

The genes encoding type I interferons (IFNs), which include fourteen IFN- α genes, one IFN- β gene, an IFN- ω (also known as IFN- α 11) gene, and a number of IFN- ω pseudogenes, are clustered on human chromosome 9p21.3. Interferon- α and - β are cytokines that are widely known to induce potent antiviral activity. They exert a variety of other biological effects, including antitumor and immunomodulatory activities and are increasingly used clinically to treat a range of malignancies, myelodysplasias and autoimmune diseases. IFN- ω is antigenically different from human IFN- α , IFN- β or IFN- γ , but is a component of natural mixtures of IFN species produced by virus-induced leukocytes or Burkitt's lymphoma cells. The type I interferon receptor (IFN- α R) interacts with IFN- α , IFN- β and IFN- ω , and seems to be a multisubunit receptor. IFN- α 7 (interferon α -7), also known as LelF J (interferon α -J) or IFN- α -J1, is a 189 amino acid secreted protein that has antiviral activity and belongs to the α / β interferon family.

REFERENCES

1. Daugherty, B., Martin-Zanca, D., Kelder, B., Collier, K., Seamans, T.C., Hotta, K. and Pestka, S. 1984. Isolation and bacterial expression of a murine α leukocyte interferon gene. *J. Interferon Res.* 4: 635-643.
2. Su, Y.H., Oakes, J.E. and Lausch, R.N. 1993. Mapping the genetic region coding for herpes simplex virus resistance to mouse interferon α / β . *J. Gen. Virol.* 74: 2325-2332.
3. Politis, A.D., Sivo, J. and Vogel, S.N. 1993. Multiple pathways of interferon-induced gene expression in murine macrophages. *J. Leukoc. Biol.* 53: 583-590.
4. Hertzog, P.J., Hwang, S.Y. and Kola, I. 1994. Role of interferons in the regulation of cell proliferation, differentiation, and development. *Mol. Reprod. Dev.* 39: 226-232.
5. Schiavoni, G., Mauri, C., Carlei, D., Belardelli, F., Pastoris, M.C. and Proietti, E. 2004. Type I IFN protects permissive macrophages from *Legionella pneumophila* infection through an IFN- γ -independent pathway. *J. Immunol.* 173: 1266-1275.
6. Chen, J., Baig, E. and Fish, E.N. 2004. Diversity and relatedness among the type I interferons. *J. Interferon Cytokine Res.* 24: 687-698.
7. Honda, K., Mizutani, T. and Taniguchi, T. 2004. Negative regulation of IFN- α / β signaling by IFN regulatory factor 2 for homeostatic development of dendritic cells. *Proc. Natl. Acad. Sci. USA* 101: 2416-2421.

CHROMOSOMAL LOCATION

Genetic locus: Ifna7 (mouse) mapping to 4 C4.

PRODUCT

IFN- α 7 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IFN- α 7 shRNA Plasmid (m): sc-146166-SH and IFN- α 7 shRNA (m) Lentiviral Particles: sc-146166-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IFN- α 7 siRNA (m) is recommended for the inhibition of IFN- α 7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IFN- α 7 gene expression knockdown using RT-PCR Primer: IFN- α 7 (m)-PR: sc-146166-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.