



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IFT80 siRNA (m): sc-146176

BACKGROUND

Intraflagellar transport is mediated by a variety of intraflagellar transport proteins (IFTs) that work in tandem to mediate ciliary and flagellar process assembly. IFT proteins are divided into two subcomplexes, A and B. IFT-A proteins are associated with retrograde transport, whereas IFT-B proteins are thought to be involved in structure because, in their absence, cilia and flagella may be truncated, or completely absent. IFT80 (intraflagellar transport protein 80 homolog) is also known as WDR56 (WD repeat-containing protein 56) and is a 777 amino acid protein that is expressed in brain, lung, liver, kidney, testis and ovary. IFT80 is localized to the cytoplasm, basal body and ciliary/flagellar axoneme. IFT80 is a component of the IFT-B complex and has WD 40 repeats, which are thought to be important for protein-protein interactions. An amino acid deletion and two missense mutations present in the gene encoding IFT80 may be associated with Jeune syndrome, an autosomally recessive chondrodysplasia that often leads to infant death due to severe bone constrictions of the rib cage which cause insufficiency of the respiratory system.

REFERENCES

1. Lin, B., White, J.T., Utleg, A.G., Wang, S., Ferguson, C., True, L.D., Vessella, R., Hood, L. and Nelson, P.S. 2003. Isolation and characterization of human and mouse WDR19, a novel WD-repeat protein exhibiting androgen-regulated expression in prostate epithelium. *Genomics* 82: 331-342.
2. Petříčková, K., Hasek, J., Benada, O. and Petříček, M. 2006. The WD-40 repeat protein PkwA of *Thermomonospora curvata* is associated with rapid growth and is localized in the tips of growing hyphae. *FEMS Microbiol. Lett.* 258: 187-193.
3. Beales, P.L., Bland, E., Tobin, J.L., Bacchelli, C., Tuysuz, B., Hill, J., Rix, S., Pearson, C.G., Kai, M., Hartley, J., Johnson, C., Irving, M., Elcioglu, N., Winey, M., Tada, M. and Scambler, P.J. 2007. IFT80, which encodes a conserved intraflagellar transport protein, is mutated in Jeune asphyxiating thoracic dystrophy. *Nat. Genet.* 39: 727-729.
4. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611177. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Huang, W., Kane, J.K. and Li, M.D. 2008. Identification and characterization of a long isoform of human IFT80, IFT80-L. *Biochem. Biophys. Res. Commun.* 373: 653-658.
6. Lee, E., Sivan-Loukianova, E., Eberl, D.F. and Kernan, M.J. 2008. An IFT-A protein is required to delimit functionally distinct zones in mechanosensory cilia. *Curr. Biol.* 18: 1899-1906.
7. Tsao, C.C. and Gorovsky, M.A. 2008. Tetrahymena IFT122A is not essential for cilia assembly but plays a role in returning IFT proteins from the ciliary tip to the cell body. *J. Cell Sci.* 121: 428-436.

CHROMOSOMAL LOCATION

Genetic locus: *Ift80* (mouse) mapping to 3 E1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

IFT80 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IFT80 shRNA Plasmid (m): sc-146176-SH and IFT80 shRNA (m) Lentiviral Particles: sc-146176-V as alternate gene silencing products.

For independent verification of IFT80 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146176A, sc-146176B and sc-146176C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IFT80 siRNA (m) is recommended for the inhibition of IFT80 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IFT80 gene expression knockdown using RT-PCR Primer: IFT80 (m)-PR: sc-146176-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.