



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IL-17RD siRNA (m): sc-146206

BACKGROUND

The interleukins (ILs) are a broad family of well characterized cytokines, primarily of hematopoietic cell origin, and are secreted by immune cells (mainly macrophages, B-cells or T-cells) that regulate a wide range of immune system functions. IL-17 and IL-17R are a cytokine receptor pair and, as such, are involved in initiating signal transduction cascades and in mediating aspects of inflammatory responses. IL-17RD (IL 17 receptor D) is also known as Sef homolog, hSef or SEF and is a 739 amino acid protein that is expressed in a variety of tissues, including ovary, breast, kidney, heart, skeletal muscle, colon, prostate and thyroid gland. IL-17RD is localized to the cytoplasm, as well as to both the Golgi apparatus membrane and to the cellular membrane as a single-pass membrane protein. IL-17RD modulates the signaling of FGFs (fibroblast growth factors) which are important for cellular proliferation, migration, differentiation and cellular survival. IL-17RD can form homomeric complexes and is thought to mediate the signaling of IL-17 by forming a heteromeric complex with IL-17R. IL-17R family members create different receptor complexes allowing interactions between different ligands. IL-17RD mRNA is detected in psoriatic skin lesions in lower concentrations than in nonlesional psoriatic skin. IL-17RD is thought to act as a tumor suppressor in humans because it is downregulated in a variety of cancers affecting epithelial cells of breast, prostate, thyroid gland and ovary.

REFERENCES

1. Yang, R.B., Ng, C.K., Wasserman, S.M., Kömüves, L.G., Gerritsen, M.E. and Topper, J.N. 2003. A novel interleukin-17 receptor-like protein identified in human umbilical vein endothelial cells antagonizes basic fibroblast growth factor-induced signaling. *J. Biol. Chem.* 278: 33232-33238.
2. Xiong, S., Zhao, Q., Rong, Z., Huang, G., Huang, Y., Chen, P., Zhang, S., Liu, L. and Chang, Z. 2003. hSef inhibits PC-12 cell differentiation by interfering with Ras-mitogen-activated protein kinase MAPK signaling. *J. Biol. Chem.* 278: 50273-50282.
3. Torii, S., Kusakabe, M., Yamamoto, T., Maekawa, M. and Nishida, E. 2004. Sef is a spatial regulator for Ras/MAP kinase signaling. *Dev. Cell* 7: 33-44.
4. Preger, E., Ziv, I., Shabtay, A., Sher, I., Tsang, M., Dawid, I.B., Altuvia, Y. and Ron, D. 2004. Alternative splicing generates an isoform of the human Sef gene with altered subcellular localization and specificity. *Proc. Natl. Acad. Sci. USA* 101: 1229-1234.
5. Zisman-Rozen, S., Fink, D., Ben-Izhak, O., Fuchs, Y., Brodski, A., Kraus, M.H., Bejar, J. and Ron, D. 2007. Downregulation of Sef, an inhibitor of receptor tyrosine kinase signaling, is common to a variety of human carcinomas. *Oncogene* 26: 6093-6098.
6. Ren, Y., Cheng, L., Rong, Z., Li, Z., Li, Y., Zhang, X., Xiong, S., Hu, J., Fu, X.Y. and Chang, Z. 2008. hSef potentiates EGF-mediated MAPK signaling through affecting EGFR trafficking and degradation. *Cell. Signal.* 20: 518-533.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 606807. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: IL17rd (mouse) mapping to 14 A3.

PRODUCT

IL-17RD siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-17RD shRNA Plasmid (m): sc-146206-SH and IL-17RD shRNA (m) Lentiviral Particles: sc-146206-V as alternate gene silencing products.

For independent verification of IL-17RD (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146206A, sc-146206B and sc-146206C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-17RD siRNA (m) is recommended for the inhibition of IL-17RD expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-17RD gene expression knockdown using RT-PCR Primer: IL-17RD (m)-PR: sc-146206-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.