



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IMMP1L siRNA (m): sc-146225

BACKGROUND

Composed of two catalytic subunits, designated IMP1 and IMP2, the mitochondrial inner membrane peptidase (IMP) complex functions in the production of active, mature mitochondrial inner membrane proteins by removing the mitochondrial targeting presequence of nuclear encoded proteins. IMMP1L (mitochondrial inner membrane protease subunit 1), also known as IMP1, is a 166 amino acid protein that belongs to the peptidase S26 family and IMP1 subfamily. Localizing to the mitochondrial inner membrane, IMMP1L participates in the assembly of the active form of Diablo, a mitochondrial protein that promotes apoptosis. The gene encoding IMMP1L maps to human chromosome 11p13 and mouse chromosome 2 E3. Chromosome 11 houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that maps to chromosome 11.

REFERENCES

- Burri, L., Strahm, Y., Hawkins, C.J., Gentle, I.E., Puryer, M.A., Verhagen, A., Callus, B., Vaux, D. and Lithgow, T. 2005. Mature DIABLO/Smac is produced by the IMP protease complex on the mitochondrial inner membrane. *Mol. Biol. Cell* 16: 2926-2933.
- Taylor, T.D., Noguchi, H., Totoki, Y., Toyoda, A., Kuroki, Y., Dewar, K., Lloyd, C., Itoh, T., Takeda, T., Kim, D.W., She, X., Barlow, K.F., Bloom, T., Bruford, E., Chang, J.L., Cuomo, C.A., Eichler, E., Fitzgerald, M.G., et al. 2006. Human chromosome 11 DNA sequence and analysis including novel gene identification. *Nature* 440: 497-500.
- Schuchman, E.H. 2007. The pathogenesis and treatment of acid sphingomyelinase-deficient Niemann-Pick disease. *J. Inherit. Metab. Dis.* 30: 654-663.
- Bhuiyan, Z.A., Momenah, T.S., Amin, A.S., Al-Khadra, A.S., Alders, M., Wilde, A.A. and Mannens, M.M. 2008. An intronic mutation leading to incomplete skipping of exon-2 in KCNQ1 rescues hearing in Jervell and Lange-Nielsen syndrome. *Prog. Biophys. Mol. Biol.* 98: 319-327.
- Mai, Y. and Gao, G. 2010. Expression of IMP1 enhances production of murine leukemia virus vector by facilitating viral genomic RNA packaging. *PLoS ONE* 5: e15881.
- Edison, N., Zuri, D., Maniv, I., Bornstein, B., Lev, T., Gottfried, Y., Kemeny, S., Garcia-Fernandez, M., Kagan, J. and Larisch, S. 2011. The IAP-antagonist ARTS initiates caspase activation upstream of cytochrome C and SMAC/Diablo. *Cell Death Differ.* 19: 356-368.
- Pluta, P., Cebula-Obrzut, B., Ehemann, V., Pluta, A., Wierzbowska, A., Piekarski, J., Bilski, A., Nejc, D., Kordek, R., Robak, T., Smolewski, P. and Jezierski, A. 2011. Correlation of Smac/DIABLO protein expression with the clinico-pathological features of breast cancer patients. *Neoplasma* 58: 430-435.
- Fullston, T., Mitchell, M., Wakefield, S. and Lane, M. 2011. Mitochondrial inhibition during preimplantation embryogenesis shifts the transcriptional profile of fetal mouse brain. *Reprod. Fertil. Dev.* 23: 691-701.

CHROMOSOMAL LOCATION

Genetic locus: *Immp1l* (mouse) mapping to 2 E3.

PRODUCT

IMMP1L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IMMP1L shRNA Plasmid (m): sc-146225-SH and IMMP1L shRNA (m) Lentiviral Particles: sc-146225-V as alternate gene silencing products.

For independent verification of IMMP1L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146225A, sc-146225B and sc-146225C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IMMP1L siRNA (m) is recommended for the inhibition of IMMP1L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IMMP1L gene expression knockdown using RT-PCR Primer: IMMP1L (m)-PR: sc-146225-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.