



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Integrin α 10 siRNA (m): sc-146246

BACKGROUND

Integrin α 10 (Integrin α -10/ β 1) is a member of the integrin α chain family that contains seven FG-GAP repeats and one VWFA domain. Integrin α 10 is single-pass type 1 membrane protein and is expressed as a hetero-dimer of an α and a β subunit. Integrin α 10 is a widely expressed protein with highest expression found in muscle and heart tissue but is also found in articular cartilage. The α 10 subunit is part of a collagen type II-binding integrin found in chondrocytes. Disruption of Integrin α 10 expression will lead to growth retardation and defects in the growth plate, and is characterized by a disturbed arrangement of chondrocytes, abnormal chondrocyte shape and reduced chondrocyte proliferation. AP-2 ϵ and Ets-1 have been shown to be involved in the regulation of Integrin α 10 transcription in chondrocytes. Integrin α 10 is upregulated in malignant melanoma cell lines.

REFERENCES

1. Lehnert, K., Ni, J., Leung, E., Gough, S., Morris, C.M., Liu, D., Wang, S.X., Langley, R. and Krissansen, G.W. 1999. The Integrin α 10 subunit: expression pattern, partial gene structure, and chromosomal localization. *Cytogenet. Cell Genet.* 87: 238-244.
2. Bengtsson, T., Camper, L., Schneller, M. and Lundgren-Akerlund, E. 2001. Characterization of the mouse integrin subunit α 10 gene and comparison with its human homologue. Genomic structure, chromosomal localization and identification of splice variants. *Matrix Biol.* 20: 565-576.
3. Gullberg, D.E. and Lundgren-Akerlund, E. 2002. Collagen-binding I domain integrins—what do they do? *Prog. Histochem. Cytochem.* 37: 3-54.
4. Bengtsson, T., Aszodi, A., Nicolae, C., Hunziker, E.B., Lundgren-Akerlund, E. and Fässler, R. 2005. Loss of α 10 β 1 Integrin expression leads to moderate dysfunction of growth plate chondrocytes. *J. Cell Sci.* 118: 929-936.
5. Wenke, A.K., Rothhammer, T., Moser, M. and Bosserhoff, A.K. 2006. Regulation of Integrin α 10 expression in chondrocytes by the transcription factors AP-2 ϵ and Ets-1. *Biochem. Biophys. Res. Commun.* 345: 495-501.
6. Farina, A., Volinia, S., Arcelli, D., Francioso, F., Desanctis, P., Zucchini, C., Pili, G., Carinci, P., Morano, D., Pittalis, M.C., Calderoni, P., Vagnoni, S. and Rizzo, N. 2006. Evidence of genetic underexpression in chorionic villi samples of euploid fetuses with increased nuchal translucency at 10-11 weeks' gestation. *Prenat. Diagn.* 26: 128-133.
7. Wenke, A.K., Kjellman, C., Lundgren-Akerlund, E., Uhlmann, C., Haass, N.K., Herlyn, M. and Bosserhoff, A.K. 2007. Expression of Integrin α 10 is induced in malignant melanoma. *Cell. Oncol.* 29: 373-386.
8. Varas, L., Ohlsson, L.B., Honeth, G., Olsson, A., Bengtsson, T., Wiberg, C., Bockermann, R., Järnum, S., Richter, J., Pennington, D., Johnstone, B., Lundgren-Akerlund, E. and Kjellman, C. 2007. α 10 Integrin expression is up-regulated on fibroblast growth factor-2-treated mesenchymal stem cells with improved chondrogenic differentiation potential. *Stem Cells Dev.* 16: 965-978.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Itga10 (mouse) mapping to 3 F2.1.

PRODUCT

Integrin α 10 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Integrin α 10 shRNA Plasmid (m): sc-146246-SH and Integrin α 10 shRNA (m) Lentiviral Particles: sc-146246-V as alternate gene silencing products.

For independent verification of Integrin α 10 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-146246A, sc-146246B and sc-146246C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Integrin α 10 siRNA (m) is recommended for the inhibition of Integrin α 10 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Integrin α 10 gene expression knockdown using RT-PCR Primer: Integrin α 10 (m)-PR: sc-146246-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.