



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



ISCA2 siRNA (m): sc-146293

BACKGROUND

ISCA2 (mitochondrial iron-sulfur cluster assembly 2 homolog, HESB-like domain-containing protein 1) is a mitochondrial protein believed to be involved in the assembly of iron and sulfur containing proteins. Iron-sulfur (Fe-S) clusters are cofactors that are essential for a wide variety of processes, including facilitation of electron transfer processes in oxidative phosphorylation, catalysis of enzymatic reactions in aconitase and dehydratases and maintenance of structural integrity in the DNA repair enzyme endonuclease III. In bacteria and eukaryotes, several new genes are implicated in the biogenesis of Fe-S cluster-containing proteins. ISCA2 is found on chromosome 14 which contains about 700 genes and 106 million base pairs and makes up about 3.5% of human cellular DNA.

REFERENCES

1. Skovran, E. and Downs, D.M. 2000. Metabolic defects caused by mutations in the *isc* gene cluster in *Salmonella enterica* serovar typhimurium: implications for thiamine synthesis. *J. Bacteriol.* 182: 3896-3903.
2. Morimoto, K., et al. 2003. A HEAT-repeats containing protein, IaiH, stabilizes the iron-sulfur cluster bound to the cyanobacterial IscA homologue, ISCA2. *J. Biochem.* 134: 211-217.
3. Cózar-Castellano, I., et al. 2004. hISCA: a protein implicated in the biogenesis of iron-sulfur clusters. *Biochim. Biophys. Acta* 1700: 179-188.
4. Berndt, C., et al. 2004. Characterization and reconstitution of a 4Fe-4S adenylyl sulfate/phosphoadenylyl sulfate reductase from *Bacillus subtilis*. *J. Biol. Chem.* 279: 7850-7855.
5. Jarrett, J.T. 2005. The novel structure and chemistry of iron-sulfur clusters in the adenosylmethionine-dependent radical enzyme biotin synthase. *Arch. Biochem. Biophys.* 433: 312-321.
6. Parikh, H., et al. 2008. Molecular correlates for maximal oxygen uptake and type 1 fibers. *Am. J. Physiol. Endocrinol. Metab.* 294: E1152-E1159.
7. Nilsson, R., et al. 2009. Discovery of genes essential for heme biosynthesis through large-scale gene expression analysis. *Cell Metab.* 10: 119-130.
8. Crooks, D.R., et al. 2010. Posttranslational stability of the heme biosynthetic enzyme ferrochelatase is dependent on iron availability and intact iron-sulfur cluster assembly machinery. *Blood* 115: 860-869.

CHROMOSOMAL LOCATION

Genetic locus: *IscA2* (mouse) mapping to 12 D1.

PRODUCT

ISCA2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ISCA2 shRNA Plasmid (m): sc-146293-SH and ISCA2 shRNA (m) Lentiviral Particles: sc-146293-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ISCA2 siRNA (m) is recommended for the inhibition of ISCA2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ISCA2 gene expression knockdown using RT-PCR Primer: ISCA2 (m)-PR: sc-146293-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.