



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



ISOC2 siRNA (m): sc-146301

BACKGROUND

ISOC2 (isochorismatase domain-containing protein 2) is a 205 amino acid protein belonging to the isochorismatase family. Isochorismatase is an enzyme that catalyzes the chemical reaction of isochorismate and water to 2,3-dihydroxy-2,3-dihydrobenzoate and pyruvate. Localized to mitochondrion and cytoplasm, ISOC2 interacts with CDKN2A and localizes to the nucleus in the presence of it. ISOC2 is ubiquitously expressed, with highest levels found in uterus, stomach and urinary tract system. Overexpressed ISOC2 inhibits the expression of CDKN2A, suggesting that this novel gene may play a role during tumor development by interacting with CDKN2A. Three isoforms are produced by alternative splicing events.

REFERENCES

1. Young, I.G. and Gibson, F. 1969. Regulation of the enzymes involved in the biosynthesis of 2,3-dihydroxybenzoic acid in *Aerobacter aerogenes* and *Escherichia coli*. *Biochim. Biophys. Acta* 177: 401-411.
2. Ruas, M. and Peters, G. 1998. The p16^{INK4a}/CDKN2A tumor suppressor and its relatives. *Biochim. Biophys. Acta* 1378: F115-F177.
3. Jacobs, J.J. and de Lange, T. 2004. Significant role for p16^{INK4a} in p53-independent telomere-directed senescence. *Curr. Biol.* 14: 2302-2308.
4. Ishikawa, A., Sasaki, M., Sato, Y., Ohira, S., Chen, M.F., Huang, S.F., Oda, K., Nimura, Y. and Nakanuma, Y. 2004. Frequent p16^{INK4a} inactivation is an early and frequent event of intraductal papillary neoplasm of the liver arising in hepatolithiasis. *Hum. Pathol.* 35: 1505-1514.
5. Ohtani, N., Yamakoshi, K., Takahashi, A. and Hara, E. 2004. The p16^{INK4a}-RB pathway: molecular link between cellular senescence and tumor suppression. *J. Med. Invest.* 51: 146-153.
6. Huang, X., Shi, Z., Wang, W., Bai, J., Chen, Z., Xu, J., Zhang, D. and Fu, S. 2007. Identification and characterization of a novel protein ISOC2 that interacts with p16^{INK4a}. *Biochem. Biophys. Res. Commun.* 361: 287-293.
7. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 612928. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/612928>

CHROMOSOMAL LOCATION

Genetic locus: Isoc2a (mouse) mapping to 7 A1.

PRODUCT

ISOC2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ISOC2 shRNA Plasmid (m): sc-146301-SH and ISOC2 shRNA (m) Lentiviral Particles: sc-146301-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ISOC2 siRNA (m) is recommended for the inhibition of ISOC2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ISOC2 gene expression knockdown using RT-PCR Primer: ISOC2 (m)-PR: sc-146301-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.