



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

JP-45 siRNA (m): sc-146331

BACKGROUND

JP-45 (junctional-face membrane protein of 45 kDa homolog), also known as JSRP1 (junctional sarcoplasmic reticulum protein 1), is a 331 amino acid sarcoplasmic and endoplasmic reticulum membrane protein. Interacting with L-type Ca^{2+} CP $\alpha 1\text{S}$, L-type Ca^{2+} CP $\beta 1\text{B}$ and calsequestrin, JP-45 may participate in the regulation of the L-type Ca^{2+} CP $\alpha 1\text{S}$ voltage-sensitive calcium channel as well as the regulation of L-type Ca^{2+} CP $\alpha 1\text{S}$ membrane targeting and activity. JP-45 may also have a role in the excitation and contraction coupling of muscle cells via interactions with key proteins present in the sarcoplasmic reticulum (SR). JP-45's interaction with SR proteins make it an important component to skeletal muscle development and maintenance. The gene encoding JP-45 maps to human chromosome 19p13.3, silencing of this gene, while having no effect on postnatal development, may result in decreased muscle strength.

REFERENCES

1. Zorzato, F., et al. 2000. Identification of a novel 45 kDa protein (JP-45) from rabbit sarcoplasmic-reticulum junctional-face membrane. *Biochem. J.* 351: 537-543.
2. Anderson, A.A., et al. 2003. The novel skeletal muscle sarcoplasmic reticulum JP-45 protein. Molecular cloning, tissue distribution, developmental expression, and interaction with $\alpha 1.1$ subunit of the voltage-gated calcium channel. *J. Biol. Chem.* 278: 39987-39992.
3. Anderson, A.A., et al. 2006. The junctional SR protein JP-45 affects the functional expression of the voltage-dependent Ca^{2+} channel Cav1.1. *J. Cell Sci.* 119: 2145-2155.
4. Ríos, E. 2006. Calcium signalling in muscle: a milestone for modulation studies. *J. Physiol.* 572: 1-2.
5. Gouadon, E., et al. 2006. A possible role of the junctional face protein JP-45 in modulating Ca^{2+} release in skeletal muscle. *J. Physiol.* 572: 269-280.
6. Delbono, O., et al. 2007. Loss of skeletal muscle strength by ablation of the sarcoplasmic reticulum protein JP-45. *Proc. Natl. Acad. Sci. USA* 104: 20108-20113.
7. Newby, P.R., et al. 2010. Follow-up of potential novel Graves' disease susceptibility loci, identified in the UK WTCCC genome-wide nonsynonymous SNP study. *Eur. J. Hum. Genet.* 18: 1021-1026.

CHROMOSOMAL LOCATION

Genetic locus: *Jsrp1* (mouse) mapping to 10 C1.

PRODUCT

JP-45 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see JP-45 shRNA Plasmid (m): sc-146331-SH and JP-45 shRNA (m) Lentiviral Particles: sc-146331-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

JP-45 siRNA (m) is recommended for the inhibition of JP-45 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

JP-45 (C-4): sc-377298 is recommended as a control antibody for monitoring of JP-45 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor JP-45 gene expression knockdown using RT-PCR Primer: JP-45 (m)-PR: sc-146331-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.