



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

JTV1 siRNA (m): sc-146335

BACKGROUND

The fidelity of protein synthesis requires efficient discrimination of amino acid substrates by aminoacyl-tRNA synthetases. Aminoacyl-tRNA synthetases function to catalyze the aminoacylation of tRNAs by their corresponding amino acids, thus linking amino acids with tRNA-contained nucleotide triplets. JTV1, also known as P38 or AIMP2 (aminoacyl tRNA synthetase complex-interacting multifunctional protein 2), is a 320 amino acid nuclear and cytoplasmic protein that is a component of the aminoacyl-tRNA synthase complex. Containing one GST C-terminal domain, JTV1 mediates ubiquitination and degradation of FBP1, a transcriptional activator of c-Myc, leading to c-Myc down-regulation, which is required for alveolar type II cell differentiation. JTV1 also participates in blocking MDM2-mediated ubiquitination, degradation of p53 and functions as a proapoptotic factor.

REFERENCES

1. Quevillon, S., Robinson, J.C., Berthonneau, E., Siatecka, M. and Mirande, M. 1999. Macromolecular assemblage of aminoacyl-tRNA synthetases: identification of protein-protein interactions and characterization of a core protein. *J. Mol. Biol.* 285: 183-195.
2. Kim, M.J., Park, B.J., Kang, Y.S., Kim, H.J., Park, J.H., Kang, J.W., Lee, S.W., Han, J.M., Lee, H.W. and Kim, S. 2003. Downregulation of FUSE-binding protein and c-Myc by tRNA synthetase cofactor p38 is required for lung cell differentiation. *Nat. Genet.* 34: 330-336.
3. Halwani, R., Cen, S., Javanbakht, H., Saadatmand, J., Kim, S., Shiba, K. and Kleiman, L. 2004. Cellular distribution of Lysyl-tRNA synthetase and its interaction with Gag during human immunodeficiency virus type 1 assembly. *J. Virol.* 78: 7553-7564.
4. Ko, H.S., von Coelln, R., Sriram, S.R., Kim, S.W., Chung, K.K., Pletnikova, O., Troncoso, J., Johnson, B., Saffary, R., Goh, E.L., Song, H., Park, B.J., Kim, M.J., Kim, S., Dawson, V.L. and Dawson, T.M. 2005. Accumulation of the authentic parkin substrate aminoacyl-tRNA synthetase cofactor, p38/JTV-1, leads to catecholaminergic cell death. *J. Neurosci.* 25: 7968-7978.
5. Han, J.M., Park, B.J., Park, S.G., Oh, Y.S., Choi, S.J., Lee, S.W., Hwang, S.K., Chang, S.H., Cho, M.H. and Kim, S. 2008. AIMP2/p38, the scaffold for the multi-tRNA synthetase complex, responds to genotoxic stresses via p53. *Proc. Natl. Acad. Sci. USA* 105: 11206-11211.
6. Choi, J.W., Um, J.Y., Kundu, J.K., Surh, Y.J. and Kim, S. 2009. Multidirectional tumor-suppressive activity of AIMP2/p38 and the enhanced susceptibility of AIMP2 heterozygous mice to carcinogenesis. *Carcinogenesis* 30: 1638-1644.
7. Choi, J.W., Kim, D.G., Park, M.C., Um, J.Y., Han, J.M., Park, S.G., Choi, E.C. and Kim, S. 2009. AIMP2 promotes TNF α -dependent apoptosis via ubiquitin-mediated degradation of TRAF2. *J. Cell Sci.* 122: 2710-2715.
8. Wang, C., Wang, D.S., Liu, B.Z., Hao, P., Liu, C., Jin, D.T., Zhong, L. and Wang, C.G. 2009. Identification of the interactions between JTV1 and NLS-RAR α *in vivo* and *in vitro*. *Sichuan Da Xue Xue Bao Yi Xue Ban* 40: 382-384, 402.

CHROMOSOMAL LOCATION

Genetic locus: Aimp2 (mouse) mapping to 5 G2.

PRODUCT

JTV1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see JTV1 shRNA Plasmid (m): sc-146335-SH and JTV1 shRNA (m) Lentiviral Particles: sc-146335-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

JTV1 siRNA (m) is recommended for the inhibition of JTV1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor JTV1 gene expression knockdown using RT-PCR Primer: JTV1 (m)-PR: sc-146335-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.