



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

KCNE1L siRNA (m): sc-146357

BACKGROUND

KCNE genes encode small, single transmembrane domain peptides that associate with pore-forming α -subunits to form K^+ channels with unique characteristics. Voltage-gated K^+ channels in the plasma membrane control the repolarization and the frequency of action potentials in neurons, muscles and other excitable cells. KCNE1L (KCNE1-like), also known as KCNE5, potassium voltage-gated channel subfamily E member 1-like protein or AMMECR2, is a 142 amino acid single-pass type I membrane protein belonging to the potassium channel KCNE family. Defects of the KCNE1L gene have been linked to the cardiac and some neurological abnormalities observed in patients with AMME (Alport syndrome, mental retardation, midface hypoplasia and elliptocytosis) contiguous gene syndrome. KCNE1L is primarily expressed in skeletal muscle, brain, placenta, spinal cord and heart.

REFERENCES

1. Piccini, M., Vitelli, F., Seri, M., Galletta, L.J., Moran, O., Bulfone, A., Banfi, S., Pober, B. and Renieri, A. 1999. KCNE1-like gene is deleted in AMME contiguous gene syndrome: identification and characterization of the human and mouse homologs. *Genomics* 60: 251-257.
2. Meloni, I., Vitelli, F., Pucci, L., Lowry, R.B., Tonlorenzi, R., Rossi, E., Ventura, M., Rizzoni, G., Kashtan, C.E., Pober, B. and Renieri, A. 2002. Alport syndrome and mental retardation: clinical and genetic dissection of the contiguous gene deletion syndrome in Xq22.3 (ATS-MR). *J. Med. Genet.* 39: 359-365.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 300328. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Hofman-Bang, J., Jespersen, T., Grunnet, M., Larsen, L.A., Andersen, P.S., Kanter, J.K., Kjeldsen, K. and Christiansen, M. 2004. Does KCNE5 play a role in long QT syndrome? *Clin. Chim. Acta* 345: 49-53.
5. Ravn, L.S., Hofman-Bang, J., Diken, U., Larsen, S.O., Jensen, G., Haunso, S., Svendsen, J.H. and Christiansen, M. 2005. Relation of 97T polymorphism in KCNE5 to risk of atrial fibrillation. *Am. J. Cardiol.* 96: 405-407.
6. Lundquist, A.L., Turner, C.L., Ballester, L.Y. and George, A.L. 2006. Expression and transcriptional control of human KCNE genes. *Genomics* 87: 119-128.
7. Ravn, L.S., Aizawa, Y., Pollevick, G.D., Hofman-Bang, J., Cordeiro, J.M., Diken, U., Jensen, G., Wu, Y., Burashnikov, E., Haunso, S., Guerschicoff, A., Hu, D., Svendsen, J.H., Christiansen, M. and Antzelevitch, C. 2008. Gain of function in IKs secondary to a mutation in KCNE5 associated with atrial fibrillation. *Heart Rhythm* 5: 427-435.

CHROMOSOMAL LOCATION

Genetic locus: Kcne1l (mouse) mapping to X F2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

KCNE1L siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KCNE1L shRNA Plasmid (m): sc-146357-SH and KCNE1L shRNA (m) Lentiviral Particles: sc-146357-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KCNE1L siRNA (m) is recommended for the inhibition of KCNE1L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KCNE1L gene expression knockdown using RT-PCR Primer: KCNE1L (m)-PR: sc-146357-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.