



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



KCNG3 siRNA (m): sc-146361

BACKGROUND

Neuronal and cardiac cells are excited by voltage-gated ion channels. Voltage-gated K⁺ channels in the plasma membrane control the repolarization and the frequency of action potentials in neurons, muscles and other excitable cells. Mutations interfering with potassium ion channels are known to cause a variety of disorders. A subunit of the potassium channel, KCNG3 (potassium voltage-gated channel subfamily G member 3) is a 436 amino acid protein that modulates channel activity and forms a heteromultimer with KV2.1, KV3.1 and KCNF1. KCNG3 contains an S4 domain that may serve as the voltage-sensor and is characterized by a series of positively charged amino acids at every third position. KCNG3 localizes to the membrane, however it must be associated with KV2.1 and possibly another partner to get inserted in the plasma membrane. KCNG3 is detected in testis, pancreas, lung, kidney, ovary, small intestine, colon, thymus, adrenal gland, spinal cord and in many parts of the brain, with the exception of the cerebellum.

REFERENCES

1. Sano, Y., Mochizuki, S., Miyake, A., Kitada, C., Inamura, K., Yokoi, H., Nozawa, K., Matsushime, H. and Furuichi, K. 2002. Molecular cloning and characterization of Kv6.3, a novel modulatory subunit for voltage-gated K⁺ channel Kv2.1. *FEBS Lett.* 512: 230-234.
2. Ottschytch, N., Raes, A., Van Hoorick, D. and Snyders, D.J. 2002. Obligatory heterotetramerization of three previously uncharacterized Kv channel α -subunits identified in the human genome. *Proc. Natl. Acad. Sci. USA* 99: 7986-7991.
3. Yan, L., Figueroa, D.J., Austin, C.P., Liu, Y., Bugianesi, R.M., Slaughter, R.S., Kaczorowski, G.J. and Kohler, M.G. 2004. Expression of voltage-gated potassium channels in human and rhesus pancreatic islets. *Diabetes* 53: 597-607.
4. Ottschytch, N., Raes, A.L., Timmermans, J.P. and Snyders, D.J. 2005. Domain analysis of Kv6.3, an electrically silent channel. *J. Physiol.* 568: 737-747.
5. Fantozzi, I., Platoshyn, O., Wong, A.H., Zhang, S., Remillard, C.V., Furtado, M.R., Petruskane, O.V. and Yuan, J.X. 2006. Bone morphogenetic protein-2 upregulates expression and function of voltage-gated K⁺ channels in human pulmonary artery smooth muscle cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 291: L993-L1004.
6. Börjesson, S.I. and Elinder, F. 2008. Structure, function, and modification of the voltage sensor in voltage-gated ion channels. *Cell Biochem. Biophys.* 52: 149-174.
7. Pischalnikova, A.V. and Sokolova, O.S. 2008. The domain and conformational organization in potassium voltage-gated ion channels. *J. Neuroimmune Pharmacol.* 4: 71-82.
8. Mederos Y Schnitzler, M., Rinne, S., Skrobek, L., Renigunta, V., Schlichthörl, G., Derst, C., Gudermann, T., Daut, J. and Preisig-Müller, R. 2009. Mutation of histidine 105 in the T1 domain of the potassium channel Kv2.1 disrupts heteromerization with Kv6.3 and Kv6.4. *J. Biol. Chem.* 284: 4695-4704.

CHROMOSOMAL LOCATION

Genetic locus: Kcng3 (mouse) mapping to 17 E4.

PRODUCT

KCNG3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KCNG3 shRNA Plasmid (m): sc-146361-SH and KCNG3 shRNA (m) Lentiviral Particles: sc-146361-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KCNG3 siRNA (m) is recommended for the inhibition of KCNG3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KCNG3 gene expression knockdown using RT-PCR Primer: KCNG3 (m)-PR: sc-146361-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.